

Habitat Restoration Program Technical Team Recommended Conservation Measures for Consideration by the BDCP Steering Committee

Introduction

This handout presents draft recommended habitat restoration conservation measures developed by the Habitat Restoration Program Technical Team (HRPTT). Conservation measures are organized into five categories—floodplain, freshwater intertidal marsh, brackish intertidal marsh, channel margin, and riparian habitat restoration conservation measures. Restored freshwater intertidal marsh as used in this handout corresponds to the tule and cattail dominated elements of the BDCP tidal freshwater emergent wetland natural community; restored riparian forest and scrub is an element of the BDCP valley riparian natural community. Shallow subtidal aquatic habitats¹ are anticipated to be restored incidentally with restoration of intertidal marshes and correspond to elements of the BDCP tidal perennial aquatic community. The HRPTT and Consultant Team have been working to estimate the extent of lands potentially suitable for physical habitat restoration in each Restoration Opportunity Area (ROA) based on topography, land use, flood control, and other considerations. Based on the restoration potentials, the extent of restoration proposed for each habitat type within each ROA will be developed and refined over the course of the BDCP planning process.

The following supplemental information is provided for each conservation measure following the conservation measure description.

Rationale. This section describes the justification for proposing the conservation measure. Rationale statements are primarily directed at identifying the covered species and ecosystem benefits that would be expected with implementing the conservation measure. The identified benefits are based on scientific literature and expert opinion as expressed by HRPTT members, as provided by experts requested to present information to the HRPTT on selected topics, and relevant expert opinion expressed in other BDCP venues (e.g., working groups and other technical teams).

Implementation timeframe. This section describes the BDCP implementation period (i.e., short-term or near-term) that is likely the most appropriate period for implementing the measure.

Implementation considerations. This section describes restoration design, management, and other relevant items that may need to be addressed by the

¹ Elevations considered suitable for shallow subtidal aquatic habitat include lands with elevations extending >0-6 feet below the intertidal zone. Lands within the shallow subtidal aquatic habitat zone may be elevated to elevations suitable for restoration of intertidal marsh habitat.

1 BDCP Implementing Entity when planning implementation of the conservation
2 measure.

3
4 **Resiliency to future change.** This section provides a qualitative assessment of
5 the likely ability of the habitat restored under the conservation measure to
6 continue to provide the desired level of covered species and ecosystem benefits
7 into the future with anticipated changes in environmental conditions with climate
8 change and sea level rise.

9
10 **Uncertainties/risks.** This section describes important uncertainties associated
11 with ability of the conservation measure to achieve the desired covered species
12 and ecosystem benefits and the ecological risks that may be associated with
13 implementing the proposed conservation measure. Important uncertainties and
14 risks are those identified in the course of HRPTT deliberations, including results
15 of coarse-level evaluations of proposed restoration actions.

16
17 **Monitoring and adaptive management considerations.** This section describes
18 monitoring and adaptive management-related elements of the conservation
19 measure, including elements of implementation that may be subject to adaptive
20 management and the types of monitoring that may be appropriate for assessing the
21 effectiveness of the restoration in achieving desired ecological benefits and for
22 informing the adaptive management process.

23
24 **Reversibility.** This section qualitatively assesses the likely ability to reverse the
25 environmental outcomes of the conservation measure, if necessary.

26
27 Attachment A, *Restoration Concept Definitions*, provides additional information
28 regarding restoration design requirements and expected ecological outcomes associated
29 with each of the habitat restoration categories.

30
31 The information described above for each of the draft proposed conservation measures
32 will be expanded upon and incorporated into appropriate sections of the BDCP
33 Conservation Strategy chapter.

34 35 36 **Floodplain Habitat Restoration Conservation Measures**

37
38 **Conservation Measure FLOO1.1: Modify and operate the Fremont Weir to**
39 **increase the frequency that the Yolo Bypass floodplain is inundated.** Within the
40 Yolo Bypass/Cache Slough Complex ROA, floodplain habitat in the Yolo Bypass would
41 be designed and operated to support the physical and biological attributes described in
42 Attachment A. To increase the frequency of inundation of floodplain habitat in the Yolo
43 Bypass, the Fremont Weir would be notched to an elevation of 17.5 feet (NAVD88) and
44 fitted with an operable gate that would allow Sacramento River water to flow into the
45 Yolo Bypass when Sacramento River stage at the weir exceeds 17.5 feet. The gates

1 would be designed and operated to provide for the efficient upstream and downstream
2 fish passage to and from the Yolo Bypass into the Sacramento River. Other design
3 elements of this measure could include:

- 4 ▪ excavation of a canal to convey water past the higher elevation natural levee of
5 the Sacramento River upstream of the Fremont Weir and past accumulated
6 sediment below the Fremont Weir to the Tule Canal;
- 7 ▪ acquisition of lands, in fee-title and through conservation or flood easements,
8 suitable for restoration of seasonally inundated aquatic habitats and for
9 accommodating future sea level rise;
- 10 ▪ grading, removal of existing berms or levees, and construction of berms or
11 levees to the extent necessary to improve the distribution (e.g., wetted area) and
12 hydrodynamic characteristics (e.g., residence times, flow ramping and recession)
13 of water moving through the Yolo Bypass, prevent stranding of covered fish
14 species, and to protect property; and
- 15 ▪ construction of a structure in the Sacramento River, if needed, in the vicinity of
16 the new weir gate to encourage the passage of juvenile salmonids migrating
17 down the Sacramento River into the Bypass.

18 The new gate in the weir would be operated to provide flows into the Bypass when
19 estimates of flow in the Sacramento River indicate that flows would be sufficient to
20 provide for inundation of the Bypass for at least 45 consecutive days between December
21 15 and April 30. Initial hydrodynamic modeling indicates that, based on historical
22 conditions, the Bypass could be inundated for a duration of at least 45 days in 48 percent
23 of years compared to current conditions in which this duration of inundation is achieved
24 in 17 percent of years. When river stage at the Fremont Weir exceeds 17.5 feet but is less
25 than the current weir elevation of 33.5 feet, the weir gate would be operated to allow a
26 mean flow of 4,000-5,000 cfs to pass into the Bypass, which would inundate
27 approximately 29,000 acres of floodplain habitat.

28
29 **Rationale:** Increasing the frequency of floodplain inundation in the Yolo Bypass
30 is expected to reduce the adverse effects of stressors related to food availability,
31 habitat availability, passage, harvest, and stranding for the covered fish species by:

- 32 ▪ creating additional spawning habitat for Sacramento splittail (Sommer et
33 al.2001a,2002, 2007, 2008, Moyle 2002, Moyle et al. 2004, Feyrer et al. 2006);
- 34 ▪ creating additional juvenile rearing habitat for Chinook salmon, Sacramento
35 splittail, and possibly steelhead (Sommer et al.2001a,b, 2002, 2007, 2008,
36 Moyle 2002, Moyle et al. 2004, Feyrer et al. 2006);
- 37 ▪ increasing the production of food for rearing salmonids, splittail, and other
38 species (Sommer et al. 2001a,b, 2002, 2007, 2008, Moyle 2002, Moyle et al.
39 2004, Feyrer et al. 2006);
- 40 ▪ increasing the availability and production of food in the Delta downstream of the
41 bypass for delta smelt, longfin smelt, and other covered species by exporting
42 organic material and phytoplankton, zooplankton, and other organisms produced

1 from the inundated floodplain into the Delta (Mitsch and Gosselink 2000, Moss
2 2007)²;

3 ■ increasing the frequency that floodplain flows transport organic carbon and
4 organisms from existing and future restored intertidal marsh at the downstream
5 end of the bypass into the Delta in support of in-Delta food production for delta
6 smelt, longfin smelt, and other covered species (Mitsch and Gosselink 2000,
7 Moss 2007)²;

8 ■ increasing the duration that the floodplain is inundated during periods that the
9 Yolo Bypass is receiving water from both the Fremont Weir and the westside
10 tributaries (e.g., Cache and Putah Creeks); and

11 ■ reducing losses of Chinook salmon, sturgeon, and other fish species to stranding
12 and illegal harvest by improving passage at the Fremont Weir.

13 Increasing the frequency and duration of inundation within the Yolo Bypass is the
14 largest opportunity for increasing inundated floodplain habitat in the North Delta.
15 The Yolo Bypass provides the only opportunity for increasing the frequency and
16 duration of inundation of a floodplain in the Planning Area without restoration of
17 historical floodplain surfaces presently in other land uses. Land use in the Yolo
18 Bypass has developed to be compatible with periodic flooding.

19
20 **Recommended Implementation Timeframe:** It is anticipated that
21 implementation of this conservation measure could be initiated in the BDCP near-
22 term implementation period.

23
24 **Implementation Considerations:** There are numerous challenges to
25 implementing this measure to improve the Yolo Bypass floodplain habitat.
26 Implementation considerations include:

27 ■ coordination with the U.S. Army Corps of Engineers and other flood control
28 agencies to allow notching, construction of an operable gate, excavation of a
29 channel, operation of the Fremont Weir, and modifications to Bypass
30 topography and flow patterns;

31 ■ coordination with the Department of Fish and Game on water management
32 affecting the Yolo Wildlife Area;

33 ■ coordination with the Yolo Basin Natural Heritage Program to ensure effective
34 implementation of conservation measures under both programs;

35 ■ ensuring that the design and management of Yolo Bypass floodplain habitats
36 would be compatible and provide synergistic species and ecosystem benefits
37 with restoration of freshwater intertidal marsh habitats in the Cache Slough
38 Complex ROA (see Conservation Measure FIMA1.1);

² Generally wetland principles support this rationale (Mitsch and Gosselink. 2000, Moss 2007), but there may be indirect effects that create complex responses as illustrated in Jassby's analysis of Bay/Delta phytoplankton production (Jassby 2008).

- 1 ▪ the need to construct levees to protect private landholdings that have not been
2 secured through conservation easements;
- 3 ▪ potential for increasing mercury methylation and resuspension and downstream
4 transport of other contaminants;
- 5 ▪ opportunities for reducing the potential adverse effects of pesticides/herbicides
6 on agricultural lands by promoting organic farming practices within the Bypass;
7 and
- 8 ▪ opportunities for providing localized floodplain inundation benefits during
9 periods when Sacramento River stage is below 17.5 feet by forcing water from
10 the Toe Drain onto adjacent lands.

11
12 **Resiliency to future changes:** This conservation measure is expected to be fairly
13 resilient to future changes in hydrology and sea levels. With changes in
14 hydrology, the period of inundation is expected to occur earlier in the year than
15 under current conditions (Cayan et al. 2006). Sea level rise would be expected to
16 reduce the extent of inundated floodplain at the south end of the bypass and result
17 in tidal emergent wetlands extending into these areas. This tidal emergent
18 wetland would produce organic carbon and organisms in support of food
19 production for covered fish species.

20
21 **Uncertainties/risks:** Methylation of mercury may occur in seasonally inundated
22 floodplains and intertidal zones, making methylmercury bioavailable to plants,
23 fish, and wildlife in and downstream of the floodplain (Alpers et al. 2006).
24 Mercury loading from Cache Creek and exposure to agricultural pesticides and
25 herbicides may adversely affect habitat productivity. Requirements and the
26 effectiveness of reducing the risk of stranding juvenile fish during floodplain
27 recession require further analysis.

28
29
30 **Monitoring and adaptive management considerations:** Implementation of this
31 conservation measure would provide opportunities to adaptively manage flows in
32 the Bypass using the new operable gate in the weir to improve food production
33 and habitat conditions for covered fish species over time based on monitoring
34 results. Results of monitoring (e.g., monitoring of phytoplankton and
35 zooplankton production relative to residence time and water depth) would help
36 identify ways to improve the design and management of floodplain habitats
37 restored in future years. Some of the monitoring considerations under various
38 bypass operations include:

- 39 ▪ extent of phytoplankton, zooplankton, and macroinvertebrate production under
40 various bypass operations;
- 41 ▪ load of organic carbon, phytoplankton, zooplankton, and macroinvertebrates
42 exported into aquatic habitats in the Delta;
- 43 ▪ effects of floodplain inundation on food production downstream of the bypass;

- 1 ▪ effects of floodplain inundation on Delta turbidity;
- 2 ▪ effects of floodplain inundation on habitat conditions for delta smelt in Cache
- 3 Slough, the Toe Drain, and other habitat use areas affected by the discharge of
- 4 water from the bypass;
- 5 ▪ levels of mercury methylation and biological uptake;
- 6 ▪ habitat use by green and white sturgeon and other covered fish species; and
- 7 ▪ growth and survival of rearing Sacramento splittail and Chinook salmon.

8 Additionally, experiments could be conducted to determine if inundating small
9 areas of the bypass floodplain during drier years by placing barriers in the Toe
10 Drain would yield tangible food and habitat benefits for covered fish species.

11
12 **Reversibility:** Flow-related effects of this conservation measure are considered
13 to be easily reversible because the BDCP Implementing Entity could choose not
14 to operate the Fremont Weir gate, thus maintaining the existing inundation
15 patterns of the Yolo Bypass. New levees and berms could permanently remove
16 farm land within the footprint of these structures if they are too costly to remove.

17
18 **Conservation Measure FLOO2.1: Coordinate with flood control agencies to**
19 **identify, fund, and implement flood control projects designed and managed to**
20 **restore and maintain floodplain, channel margin, freshwater intertidal, and**
21 **transitional grassland habitats.** The BDCP Implementing Entity would coordinate
22 flood control planning with the Central Valley Flood Protection Board, California
23 Department of Water Resources (DWR), and U.S. Army Corps of Engineers to identify
24 opportunities for creating new inundated floodplain habitats associated with flood control
25 projects within the Planning Area. These coordination activities would include
26 participation in development of the Central Valley Flood Protection Control Plan
27 elements related to flood control planning within the Planning Area and upstream of the
28 Planning Area where such projects may affect implementation of the BDCP Conservation
29 Strategy. The BDCP Implementing Entity would coordinate with flood control agencies
30 to identify cost sharing opportunities for creating flood conveyance features that would
31 provide joint flood control and habitat benefits for BDCP covered species through cost-
32 sharing agreements as appropriate.

33
34 Specific opportunities to be evaluated by the BDCP Implementing Entity include
35 assessing feasibility for constructing a new flood bypass channel along the east side of
36 the Sacramento River Deep Water Ship Channel and setting back levees along the San
37 Joaquin River downstream of Vernalis.

38 39 **Deep Water Ship Channel Flood Bypass**

40
41 This concept is to create a new flood bypass in the Steamboat Slough ROA east of the
42 Sacramento Deep Water Ship Channel and west of the Sacramento River that could
43 reduce flood risks to Clarksburg and the Pocket Area of Sacramento and reduce flood
44 pressures along downstream levees to Rio Vista. If implemented, the bypass would be

1 designed and operated to provide seasonally inundated floodplain habitat for periods of at
2 least up to 45 days from late-winter through spring during years when sufficient water is
3 available in the Sacramento River for this purpose. Restored floodplain habitat within the
4 bypass would be designed and operated to support the physical and biological attributes
5 described in Attachment A.

6
7 Design elements of this measure could include:

- 8 ▪ acquisition of lands in fee-title or through conservation easements suitable for
9 restoration of seasonally inundated floodplain habitats and for accommodating
10 future sea level rise;
- 11 ▪ construction of a new levee east of the Sacramento Deep Water Ship Channel to
12 contain bypass flows between the new levee and the existing east levee of the
13 Deep Water Ship Channel (the bypass width would be relatively narrow [an
14 estimated 1,000-2,000 feet] to minimize impacts on existing land uses and still
15 provide substantial benefits to covered species);
- 16 ▪ construction of an operable weir along the west levee Sacramento River upstream
17 of Freeport designed to pass flows into the bypass and to provide for passage of
18 fish upstream and downstream of the weir;
- 19 ▪ modify the landform within the bypass to prevent stranding of covered fish
20 species.
- 21 ▪ removing levees at the south end of the bypass to provide flow connectivity with
22 the Delta; and
- 23 ▪ potentially discontinuing farming within the bypass if the bypass is designed with
24 sufficient flood capacity to provide for the natural establishment and growth of
25 riparian vegetation on the floodplain surface to provide structural and
26 hydrodynamic complexity (the bypass width likely would be too narrow to
27 provide for both farming and the desired level of riparian habitat-associated
28 benefits).

29 Preliminary assessments of this concept indicate that, based on flows recorded at Freeport
30 from 1984-2007, a weir invert elevation of 6 feet in the vicinity of Freeport would allow
31 at least 3,000 cfs to inundate the floodplain for at least 45 consecutive days in 48 percent
32 of the years . The extent of inundated floodplain would be determined by the width of
33 the bypass, but would be expected to range between 2,000 and 5,000 acres.

34 35 **San Joaquin River Setback Levees—Vernalis to Mossdale**

36
37 Located within the South Delta ROA, this concept is to expand the flood capacity of the
38 existing constricted flood control channel downstream of Vernalis to Mossdale by setting
39 back levees along the San Joaquin River to expand the floodplain to allow flood waters to
40 attenuate, improving access of juvenile fish, such as Chinook salmon and steelhead, to
41 seasonally inundated floodplain habitat, and reducing flood risk to properties upstream
42 and downstream. If implemented, restored floodplain habitat along the San Joaquin
43 River would be designed and operated to support the physical and biological attributes

1 described in Attachment A. Implementation would require acquisition of lands in fee-
2 title or through conservation easements within the footprint of the expanded floodway
3 and levees.

4
5 Floodplain habitat would be restored by setting back levees along the San Joaquin River
6 and removing all or large sections of the existing levees. The extent that levees would be
7 set back and the extent of floodplain restored would primarily be dependent on the extent
8 of restored floodplain that could be inundated under [redacted] year flood events as modeled for
9 hydrological conditions expected with climate change. Initial hydrodynamic modeling
10 under existing hydrologic conditions suggests that, on average, new floodplain habitat
11 areas could be inundated for at least 30 consecutive days from late winter to early spring
12 on average once every 5.5 years (i.e., 18% of years). The new floodplain area would be
13 contoured, if needed, to reduce and avoid the potential for stranding of juvenile and adult
14 fish following inundation events.

15 The channel within the restored floodplain reach would be modified where practicable to
16 create backwater (i.e., low velocity) habitat areas designed to provide spawning habitat
17 for splittail and rearing habitat for splittail and salmonids. Within the restored floodplain,
18 farming potentially would be discontinued and riparian vegetation would be allowed to
19 naturally establish and the channel would be allowed to meander between the new levees
20 through the natural processes of erosion and sedimentation (the width of setback levees
21 likely would be too narrow to provide for both farming and the desired level of riparian
22 habitat-associated benefits).

23 **.San Joaquin River Setback Levees—Mossdale to French Camp Slough**

24
25 Located within the South Delta ROA, this concept is to increase seasonally inundated
26 floodplain habitat and expand the flood capacity of the existing flood control channel
27 downstream of Mossdale to French Camp Slough by setting back levees along the San
28 Joaquin River. Restored floodplain habitat would be designed and operated to support
29 the physical and biological attributes described in Attachment A. Implementation would
30 require acquisition of lands in fee-title or through conservation easements within the
31 footprint of the expanded floodway and levees.

32
33 Floodplain habitat would be restored by setting back levees along the San Joaquin River
34 and removing all or large sections of the existing levees. The extent to which levees
35 would be setback and the extent of floodplain habitat restored would primarily be
36 dependent on the extent of restored floodplain that could be inundated under [redacted] year
37 flood events as modeled for hydrological conditions expected with climate change and
38 land surface elevations. The new floodplain area would be contoured, if needed, to
39 reduce and avoid the potential for stranding of juvenile and adult fish following
40 inundation events. Ground surface elevations along tidal reaches may need to be elevated
41 to allow natural establishment of tidal freshwater wetland and riparian habitat.

42 The channel within the restored floodplain reach would be modified where practicable to
43 create lower velocity backwater habitat areas designed to provide spawning habitat for
44 splittail and rearing habitat for splittail and salmonids. Within the restored floodplain,
45 farming potentially would be discontinued and riparian vegetation would be allowed to

1 naturally establish and the channel would be allowed to meander between the new levees
2 through the natural processes of erosion and sedimentation (the width of setback levees
3 likely would be too narrow to provide for both farming and the desired level of riparian
4 habitat-associated benefits).

5
6 **Rationale:** Flood control agencies are currently planning modifications to the
7 existing Central Valley flood control system, which provides an opportunity for
8 the BDCP Implementing Entity to coordinate with these agencies to help ensure
9 that proposed modifications provide habitat and ecosystem benefits for covered
10 species.

11
12 Increasing the extent of floodplain habitat by creating the Deep Water Ship
13 Channel bypass is expected to reduce the adverse effects of stressors related to
14 food and habitat availability for the covered fish species by:

- 15 ■ creating additional spawning habitat for Sacramento splittail by expanding
16 access to floodplain habitat area and providing in-channel spawning habitat by
17 creating backwaters (Sommer et al.2001a, 2002, 2007, 2008, Moyle 2002,
18 Moyle et al. 2004, Feyrer et al. 2006);
- 19 ■ creating additional rearing habitat for San Joaquin River runs of Chinook
20 salmon, Sacramento splittail, and possibly steelhead (Sommer et al.2001a,b,
21 2002, 2007, 2008, Moyle 2002, Moyle et al. 2004, Feyrer et al. 2006);
- 22 ■ increasing the production of food for rearing salmonids, splittail, and other
23 species (Sommer et al. 2001a,b, 2002, 2007, 2008, Moyle 2002, Moyle et al.
24 2004, Feyrer et al. 2006);
- 25 ■ naturally establishing freshwater intertidal marsh at suitable elevations within
26 the bypass as a result of restoring tidal connectivity that will produce organic
27 carbon and food in support of aquatic food web processes;
- 28 ■ increasing the availability and production of food in Delta channels downstream
29 of restored floodplain habitat for delta smelt, longfin smelt, and other covered
30 species by exporting organic material and phytoplankton, zooplankton, and
31 other organisms produced from the inundated floodplain into the Delta (Mitsch
32 and Gosselink 2000, Moss 2007)².

33
34 In addition to providing benefits for the covered fish species, riparian habitats
35 established within the new floodplain would substantially increase valley
36 elderberry longhorn beetle habitat and Swainson's hawk nesting habitat.

37
38 Increasing the extent of floodplain habitat by setting back levees along the San
39 Joaquin River is expected to reduce the adverse effects of stressors related to food
40 and habitat availability for the covered fish species by:

- 41 ■ creating additional spawning habitat for Sacramento splittail by expanding
42 floodplain habitat area and providing in-channel spawning habitat by creating
43 backwaters Sommer et al.2001a,2002, 2007, 2008, Moyle 2002, Moyle et al.

- 1 2004, Feyrer et al. 2006)
- 2 ▪ creating additional rearing habitat for San Joaquin River runs of Chinook
- 3 salmon, Sacramento splittail, and possibly steelhead (Sommer et al.2001a,b,
- 4 2002, 2007, 2008, Moyle 2002, Moyle et al. 2004, Feyrer et al. 2006);
- 5 ▪ increasing the production of food for rearing salmonids, splittail, and other
- 6 species (Sommer et al. 2001a,b, 2002, 2007, 2008, Moyle 2002, Moyle et al.
- 7 2004, Feyrer et al. 2006);
- 8 ▪ increasing the availability and production of food in Delta channels downstream
- 9 of restored floodplain habitat for delta smelt, longfin smelt, and other covered
- 10 species by exporting organic material and phytoplankton, zooplankton, and
- 11 other organisms produced from the inundated floodplain into Delta channels
- 12 (Mitsch and Gosselink 2000, Moss 2007)²; and
- 13 ▪ increasing habitat complexity by allowing the natural establishment and growth
- 14 of woody riparian vegetation that will provide inputs of large woody debris into
- 15 the river channel and provide overhead cover.

16

17 Preliminary results of hydrodynamic modeling of dual conveyance scenarios

18 indicates that restoring habitat along the San Joaquin River between Vernalis and

19 French Camp Slough would minimize the potential for entrainment of organic

20 carbon, food, and fish produced from the restored habitat into the south Delta SWP

21 and CVP pumping facilities relative to other locations where floodplain habitats

22 could be restored in the south Delta.

23

24 In addition to providing benefits for the covered fish species, riparian habitats

25 established within the new floodplain habitat along the San Joaquin River would

26 substantially increase habitat for Swainson's hawk, riparian brush rabbit, valley

27 elderberry longhorn beetle, delta button celery, and delta tule pea.

28

29 **Recommended Implementation Timeframe:** It is anticipated that the restoration

30 actions described under this conservation measure would be implemented in the

31 BDCP long-term implementation period to provide adequate time for completion

32 of necessary interagency coordination and planning processes with local

33 landowners. This conservation measure would not be implemented until after

34 completion of the around-Delta conveyance facilities to minimize adverse effects

35 of South Delta SWP and CVP pumping operations on the functions of the restored

36 habitat. Interagency coordination and planning with local landowners, however,

37 could be initiated in the near-term implementation period.

38

39 **Implementation Considerations:** Implementation considerations include:

- 40 ▪ coordination with the U.S. Army Corps of Engineers and other flood control
- 41 agencies to allow for 1) the removal of flood control levees and the construction
- 42 of new flood control levees setback from San Joaquin River and 2) creating a
- 43 new weir on the west side of the Sacramento River and using the east Deep
- 44 Water Ship Channel levee as the west levee of the new flood bypass;

- 1 ▪ ensuring that designs would be compatible and provide synergistic species and
2 ecosystem benefits with restoration of floodplain habitats along Old River or
3 Middle River (see Conservation Measure FLOO2.2) and freshwater tidal marsh
4 habitats in the Steamboat Slough ROA (see Conservation Measure IFMA1.5)
5 and South Delta ROA (see Conservation Measure FIMA1.4);
- 6 ▪ potential for increasing mercury methylation and resuspension and downstream
7 transport of other contaminants;
- 8 ▪ potential for short-term mobilization of toxic compounds from newly inundated
9 agricultural lands;
- 10 ▪ potential for aggravating low DO in the Stockton Deep Water Ship Channel if
11 late floods produce large amounts of algae or decaying organic material that are
12 transported into the Ship Channel;
- 13 ▪ opportunities for increasing the frequency of inundation of the restored
14 floodplain in future years if changes in upstream operations increase San
15 Joaquin River flows entering the Delta; and
- 16 ▪ potential for increased inundation frequency and duration with future changes in
17 hydrology resulting from climate change.

18
19 **Resiliency to future changes:** Floodplain and freshwater tidal marsh actions
20 described in this conservation measure are expected to be fairly resilient to future
21 changes in hydrology and sea levels. With changes in hydrology, the frequency
22 of inundation would be expected to increase and period of inundation could be
23 expected to occur earlier in winter year than under current conditions (Cayan et al.
24 2006). Sea level rise could reduce the extent of inundated floodplain in
25 downstream restored habitat area as sea level rises. The lost floodplain habitat,
26 however, would be expected to develop as tidal marsh, which would produce
27 organic carbon and organisms in support of food production for covered fish
28 species. Restored tidal marsh upstream of Stockton would be expected to
29 establish further upstream in the floodplain as sea level rises.

30
31 **Uncertainties/risks:** Methylation of mercury may occur in seasonally inundated
32 floodplains and intertidal zones, making methylmercury bioavailable to plants,
33 fish, and wildlife in and downstream of the floodplain (Alpers et al. 2006).
34 Exposure to agricultural pesticides and herbicides may impact habitat productivity
35 in the first few periods that the restored floodplain is inundated. Requirements
36 and the effectiveness of reducing the risk of stranding juvenile fish during
37 floodplain recession require further analysis.

38
39 **Monitoring and adaptive management considerations:** Opportunities for
40 adaptive management are associated with assessing the effectiveness of in-
41 channel backwater and seasonal floodplain habitat restoration designs and the
42 ability of native riparian vegetation to successfully establish on new floodplain
43 surfaces and along the channels. Monitoring the establishment of riparian
44 vegetation would provide information necessary for determining the need to

1 control the establishment of non-native vegetation or plant native vegetation to
2 promote development of native riparian forest and scrub habitats. Monitoring of
3 restored floodplain habitats would also provide information that would be useful
4 in restoring floodplains in other locations. Some of the monitoring considerations
5 include:

- 6 ▪ phytoplankton and zooplankton production on the inundated floodplain and
7 changes in in-channel phytoplankton and zooplankton production associated
8 with increasing the complexity of in-channel habitat;
- 9 ▪ load of organic carbon, phytoplankton, zooplankton, and macroinvertebrates
10 exported into aquatic habitat in the Delta;
- 11 ▪ effects of floodplain inundation on food production and water quality in
12 downstream areas;
- 13 ▪ effects of floodplain inundation on Delta turbidity;
- 14 ▪ habitat use by green and white sturgeon, salmon, steelhead, and other covered
15 fish;
- 16 ▪ levels of mercury methylation and resuspension of contaminants, and
17 biological uptake;
- 18 ▪ covered fish species use of restored backwaters; and
- 19 ▪ growth and survival of rearing Sacramento splittail and Chinook salmon.

20
21 **Reversibility:** The restoration actions described under this conservation measure
22 would be very difficult to reverse because of the high capital costs associated with
23 construction of new levees and the removal of existing levees.
24

25 **Conservation Measure FLOO2.2: Restore between [redacted] and [redacted] acres of inundated**
26 **floodplain habitat in the South Delta Restoration Opportunity Area.** Within the
27 South Delta ROA, inundated floodplain habitat would be restored on Fabian Tract along
28 Old River or on Union Island and Upper Roberts Island along Middle River, depending
29 on which river corridor is not used for through-Delta conveyance. Restored floodplain
30 habitat would be designed and operated to support the physical and biological attributes
31 described in Attachment A.
32

33 Design elements of this conservation measure could include:

- 34 ▪ acquisition of lands in fee-title or through conservation easements suitable for
35 restoration of intertidal and subtidal habitats and for accommodating future sea
36 level rise;
- 37 ▪ setting back levees along the selected river corridor and removing the existing
38 levees or large sections of the existing levees;
- 39 ▪ discontinuing farming within the setback levees and allowing riparian vegetation
40 to naturally establish on the floodplain; and

- 1 ▪ re-contouring the restored floodplain surface, if needed, to avoid potential for
2 stranding of juvenile and adult fish following inundation events.

3
4 **Rationale:** Increasing the extent of floodplain habitat is expected to reduce the
5 adverse effects of stressors related to food and habitat availability for the covered
6 fish species by:

- 7 ▪ creating additional spawning habitat for Sacramento splittail by expanding
8 floodplain habitat area (Sommer et al.2001a, 2002, 2007, 2008, Moyle 2002,
9 Moyle et al. 2004, Feyrer et al. 2006);
- 10 ▪ creating additional rearing habitat for Sacramento splittail, runs of Chinook
11 salmon from the San Joaquin River and other eastside tributaries, and possibly
12 steelhead (Sommer et al.2001a,b, 2002, 2007, 2008, Moyle 2002, Moyle et al.
13 2004, Feyrer et al. 2006);
- 14 ▪ increasing the production of food for rearing salmonids, splittail, and other
15 species (Sommer et al. 2001a,b, 2002, 2007, 2008, Moyle 2002, Moyle et al.
16 2004, Feyrer et al. 2006);
- 17 ▪ increasing the availability and production of food in the Delta downstream of
18 restored floodplain habitat for delta smelt, longfin smelt, and other covered
19 species by exporting organic material and phytoplankton, zooplankton, and
20 other organisms produced from the inundated floodplain into the Delta (Mitsch
21 and Gosselink 2000, Moss 2007)²; and
- 22 ▪ increasing hydrodynamic and structural complexity within the channel by
23 allowing the natural establishment and growth of woody riparian vegetation that
24 would provide inputs of large woody debris into the river channel and provide
25 overhead cover.

26 Improving in-channel habitat complexity along the Old or Middle River corridors
27 would be expected to reduce the predation risk to covered fish species and improve
28 connectivity between San Joaquin River habitats and Delta habitats for passage of
29 juvenile salmonids outmigrating from the San Joaquin River and eastside
30 tributaries.

31 In addition to providing benefits for the covered fish species, restored riparian
32 habitats associated with creating new floodplain habitat in the South Delta ROA
33 would substantially increase habitat for Swainson's hawk, riparian brush rabbit,
34 valley elderberry longhorn beetle, delta button celery, and delta tule pea.

35
36 **Recommended Implementation Timeframe:** This conservation measure would
37 not be implemented until after completion of the around-Delta conveyance
38 facilities to minimize adverse affects of South Delta SWP/CVP pumping
39 operations on the functions of the restored habitat. Restoration planning and
40 design could be initiated in the near-term implementation period.

41
42 **Implementation Considerations:** Implementation considerations include:

- 1 ▪ selecting the location for floodplain restoration (Fabian Tract, Union Island, or
2 Middle Roberts Island) is dependent on the though-Delta corridor (i.e., Old
3 River or Middle River) selected for dual operations and therefore the relative
4 influence of South Delta SWP/CVP pumping operations on the restored habitat;
- 5 ▪ coordination with the Department of Water Resources and local reclamation
6 districts to allow for the removal of flood control levees and the construction of
7 new flood control levees setback from the selected river corridor;
- 8 ▪ ensuring that designs would be compatible and provide synergistic species and
9 ecosystem benefits with restoration of floodplain habitats along the San Joaquin
10 River (see Conservation Measure FLOO2.1) and freshwater intertidal marsh
11 habitats in the South Delta ROA (see Conservation Measure FIMA1.4);
- 12 ▪ potential for increasing mercury methylation;
- 13 ▪ potential for short-term mobilization of toxic compounds from newly inundated
14 lands;
- 15 ▪ opportunities for increasing the frequency of inundation of the restored
16 floodplain in future years if changes in upstream operations increase San
17 Joaquin River flows entering the Delta; and
- 18 ▪ potential for increased inundation frequency with future changes in hydrology
19 resulting from climate change.

20
21 **Resiliency to future changes:** This conservation measure is expected to be
22 somewhat resilient to future changes in the hydrograph and sea level. With
23 changes in the hydrograph, the frequency of inundation would be expected to
24 increase and inundation could occur earlier in the year than under current
25 conditions (Cayan et al. 2006). Sea level rise could reduce the extent of inundated
26 floodplain in downstream restoration areas. The floodplain habitat inundated by
27 sea level rise, however, would be expected to develop into tidal marsh, which
28 would produce organic carbon and organisms in support of food production for
29 covered fish species.

30
31 **Uncertainties/risks:** Methylation of mercury may occur in seasonally inundated
32 floodplains and intertidal zones, making methylmercury bioavailable to plants,
33 fish, and wildlife in and downstream of the floodplain (Alpers et al. 2006).
34 Exposure to residual agricultural pesticides and herbicides may impact habitat
35 productivity in the first few periods that the restored floodplain is inundated.

36
37 **Monitoring and adaptive management considerations:** Opportunities for
38 adaptive management are related to assessing the effectiveness of restored
39 floodplain to develop as functional habitat for covered species and to produce
40 food and organic material in support of food web processes. Adaptive
41 management considerations include assessing the need for further actions to
42 improve species benefits if indicated through monitoring (e.g., control of non-
43 native fish predators if survival of outmigrating salmonids using the corridor is

1 not improved). Monitoring the establishment of riparian vegetation on the
2 restored floodplains and along the channel would also provide information useful
3 to restoring floodplains in other locations. Some of the monitoring considerations
4 include:

- 5 ▪ phytoplankton and zooplankton production on the inundated floodplain and
6 changes in in-channel phytoplankton and zooplankton production associated
7 with increasing the complexity of in-channel habitat;
- 8 ▪ load of organic carbon, phytoplankton, zooplankton, and macroinvertebrates
9 exported into aquatic habitat in the Delta;
- 10 ▪ natural establishment and growth of riparian vegetation;
- 11 ▪ effects of floodplain inundation on food production and water quality in
12 downstream areas;
- 13 ▪ effects of floodplain inundation of Delta turbidity;
- 14 ▪ habitat use by green and white sturgeon, salmon, and other covered fish
15 species;
- 16 ▪ levels of mercury methylation and biological uptake; and
- 17 ▪ growth and survival of rearing Sacramento splittail and Chinook salmon.

18

19 **Reversibility:** This conservation measure would be difficult to reverse because
20 of the high capital costs associated with construction of new levees and the
21 removal of existing levees.

22

23

24

Freshwater Intertidal Marsh Habitat Restoration Conservation Measures

25

26

27

**Conservation Measure FIMA1.1. Restore a mosaic of [redacted] to [redacted] acres of freshwater
intertidal marsh, shallow subtidal aquatic, and transitional grassland habitat within
the Yolo Bypass/Cache Slough Complex Restoration Opportunity Area.** Restored
freshwater intertidal marsh and shallow subtidal aquatic habitats would be designed to
support the physical and biological attributes described in Attachment A. The mosaic of
habitats would include at least [redacted] acres of freshwater intertidal marsh habitat. Areas
suitable for restoration include, but are not limited to, Haas Slough, Hastings Cut,
Lindsey Slough, Barker Slough, Calhoun Cut, Liberty Island, Little Holland, the
Westlands property, Shag Slough, Little Egbert Tract, and Prospect Island. Design
elements of this conservation measure could include:

37

38

39

- acquisition of lands in fee-title or through conservation easements suitable for
restoration of intertidal and subtidal habitats and for accommodating future sea
level rise;

- 1 ▪ breaching and setting back levees to provide for tidal exchange with restored
- 2 habitats;
- 3 ▪ modifying ditches and cuts to create a dendritic pattern of tidal channels;
- 4 ▪ restoring stream functions of erosion and sedimentation (e.g., Ulatis Flood
- 5 Control channel) to improve spawning conditions for delta smelt and other fish
- 6 and macroinvertebrates; and
- 7 ▪ planting tules to raise ground surface elevations suitable for tidal marsh
- 8 restoration on subsided lands (e.g., Little Egbert Tract).
- 9

10 **Rationale:** Restoring freshwater intertidal marsh and shallow subtidal aquatic

11 habitats within the Cache Slough Complex is expected to reduce the adverse

12 effects of stressors related to food availability and habitat availability for the

13 covered fish species by:

- 14 ▪ increasing rearing habitat area for Chinook salmon, Sacramento splittail, and
- 15 possibly steelhead (Healey 2001, Brown 2003);
- 16 ▪ increasing the production of food for rearing salmonids, splittail, and other
- 17 species (Kjelson et al. 1982, Siegel 2007);
- 18 ▪ increasing the availability and production of food in the Delta downstream of
- 19 Rio Vista by exporting organic material from the marsh plain and
- 20 phytoplankton, zooplankton, and other organisms produced in intertidal
- 21 channels into the Delta (Siegel 2007);
- 22 ▪ locally providing areas of cool water refugia for delta smelt (C. Enright pers.
- 23 comm.);
- 24 ▪ increasing the extent of habitat available for colonization by Mason’s lilaeopsis;
- 25 and
- 26 ▪ increasing the extent of habitat for giant garter snake, California black rail, and
- 27 tricolored blackbird.
- 28

29 Additionally, the Cache Slough Complex encompasses a substantial area of land

30 with elevations suitable for freshwater tidal marsh restoration that would involve

31 few impacts on infrastructure or permanent crops relative to other areas of the

32 north Delta.

33

34 **Recommended Implementation Timeframe:** It is anticipated that

35 implementation of this conservation measure could be initiated in the BDCP near-

36 term implementation period.

37

38 **Implementation Considerations:** Implementation considerations include:

- 39 ▪ the need to coordinate with the Solano County HCP to ensure effective
- 40 implementation of conservation measures under both programs;
- 41 ▪ feasibility for subsidence reversal using tule plantings or other techniques to

- 1 raise ground surface elevations before breaching levees;
- 2 ▪ ensuring compatibility with flood control functions of the Yolo Bypass;
- 3 ▪ ensuring that designs would be compatible and provide synergistic species
- 4 and ecosystem benefits with proposed restoration of floodplain habitats in
- 5 the Yolo Bypass as described under Conservation Measure FLOO1.1;
- 6 ▪ the need to coordinate with land owners and other conservation planning
- 7 efforts;
- 8 ▪ the need to incorporate design features and management strategies to
- 9 preclude or minimize the establishment of *Egeria* and other undesirable
- 10 non-native species;
- 11 ▪ the need to incorporate design features that will promote the natural
- 12 establishment of marsh-associated covered plant species;
- 13 ▪ consideration for the effects of restoration-induced dampening of the tidal
- 14 range on subsequent marsh restoration designs;
- 15 ▪ potential for increasing mercury methylation and resuspension of
- 16 contaminants;
- 17 ▪ locating and designing levee breaches to maximize the development of
- 18 intertidal marsh and minimize hydrodynamic conditions that favor non-
- 19 native predatory fish;
- 20 ▪ determining the appropriate allowable land uses and management activities
- 21 on transitional grasslands conserved to accommodate future sea level rise;
- 22 and
- 23 ▪ the need to address the likely adverse effects of the Barker Slough Pumping
- 24 Plant intake on entrainment of food produced from and fish inhabiting
- 25 restored marshes before restoring habitats south of Lindsey Slough.

26
27 **Resiliency to future changes:** This conservation measure is expected to be
28 resilient to future changes in hydrology and sea levels. Conserving higher
29 elevation transitional grassland habitat along the margins of restored intertidal
30 marsh would provide sufficient lands to accommodate the upslope establishment
31 of intertidal marsh as sea level rises.

32
33 **Uncertainties/risks:** Restoration of subtidal aquatic habitats could result in
34 infestation by non-native submerged aquatic vegetation and increase the
35 abundance of non-native predators or vulnerability of covered fish species to
36 predation. Methylation of mercury may occur in intertidal zones, making
37 methylmercury bioavailable to plants, fish, and wildlife in and downstream of
38 restored marshes (Alpers et al. 2006). Altering habitat conditions in this area
39 could potentially adversely affect delta smelt spawning in this area if salinity
40 gradients, turbidity, or temperature conditions that support delta smelt habitat are
41 degraded as a result of restoration actions. Additionally, there could be a short-

1 term risk associated with mobilizing pesticides, herbicides, and other
2 contaminants into the Delta following initial introduction of tidal flow onto
3 agricultural lands.

4
5 **Monitoring and adaptive management considerations:** Opportunities for
6 adaptive management are related to assessing the effectiveness of restored
7 marshes and adjacent shallow subtidal habitats to develop as functional covered
8 species habitats and to produce food and organic carbon in support of food web
9 processes. Results of monitoring the development of early marsh restorations
10 would help inform improvements in the design and management of subsequent
11 marsh restoration projects. Results of monitoring early restorations could also be
12 used to develop cost effective management techniques, if needed, to control the
13 establishment of non-native species in restored marshes. Some of the monitoring
14 considerations include:

- 15 ■ type and extent of use by covered fishes;
- 16 ■ extent of phytoplankton, zooplankton, and macroinvertebrate production in
17 marsh channels;
- 18 ■ load of organic carbon, phytoplankton, zooplankton, and macroinvertebrates
19 produced in emergent marshes and subsequently exported into the Delta;
- 20 ■ extent of native vegetation relative to non-native vegetation on the marsh plain;
- 21 ■ extent of native aquatic vegetation relative to non-native aquatic vegetation;
- 22 ■ growth and survival of rearing Sacramento splittail and Chinook salmon in
23 shallow subtidal aquatic habitats;
- 24 ■ the establishment of habitat conditions suitable for the natural establishment of
25 marsh-associated covered plant species; and
- 26 ■ levels of mercury methylation and biological uptake.

27
28 **Reversibility:** This conservation measure would be difficult to reverse because it
29 would require re-construction of levees to re-isolate restored habitat areas from
30 tidal flow and pumping to remove water from reclaimed habitat areas.

31
32 **Conservation Measure FIMA1.2: Restore a mosaic of [redacted] to [redacted] acres of freshwater**
33 **intertidal marsh, shallow subtidal aquatic, and transitional habitat within the**
34 **Cosumnes/Mokelumne ROA.** Restored freshwater intertidal marsh and shallow
35 subtidal aquatic habitats would be designed to support the physical and biological
36 attributes described in Attachment A. The mosaic of habitats would include at least [redacted]
37 acres of freshwater intertidal marsh habitat. Areas suitable for restoration include
38 McCormack-Williamson Tract, New Hope Tract, Canal Ranch Tract, Bract Tract,
39 Terminous Tract north of State Highway 12, and lands adjoining Snodgrass Slough,
40 South Stone Lake, and Lost Slough. Design elements of this conservation measure could
41 include:

- 1 ▪ acquisition of lands in fee-title or through conservation easements suitable for
- 2 restoration of intertidal and subtidal habitats and for accommodating future sea
- 3 level rise;
- 4 ▪ constructing levees to isolate deeply subsided lands and protect private
- 5 property;
- 6 ▪ planting tules or placing fill material to raise elevations of shallowly subsided
- 7 lands,
- 8 ▪ creating channels to promote the development of tidal channels; and
- 9 ▪ breaching levees to reintroduce tidal exchange to currently leveed lands.

10 If the eastern alignment of an around-Delta aqueduct is constructed, the aqueduct levees
11 may be incorporated into the design of intertidal emergent wetland restoration.

12

13 **Rationale:** Restoring freshwater intertidal marsh and shallow subtidal aquatic
14 habitats within the Cosumnes/Mokelumne River ROA is expected to reduce the
15 adverse effects of stressors related to food and habitat availability for the covered
16 fish species by:

- 17 ▪ increasing rearing habitat area for Sacramento splittail and Cosumnes and
- 18 Mokelumne River fall-run Chinook salmon and possibly steelhead (Healey
- 19 2001, Brown 2003);
- 20 ▪ increasing the production of food for rearing salmonids, splittail, and other
- 21 species migrating to and from the Cosumnes and Mokelumne Rivers (Kjelson et
- 22 al. 1982, Siegel 2007);
- 23 ▪ increasing the availability and production of food in the east and central Delta
- 24 by exporting organic material from the marsh plain and phytoplankton,
- 25 zooplankton, and other organisms produced in intertidal channels into the Delta
- 26 (Siegel 2007);
- 27 ▪ locally providing areas of cool water refugia for delta smelt (C. Enright pers.
- 28 comm.);
- 29 ▪ increasing the extent of habitat available for colonization by Mason's lilaeopsis,
- 30 and
- 31 ▪ increasing the extent of habitat for giant garter snake, California black rail, and
- 32 tricolored blackbird.

33

34 **Recommended Implementation Timeframe:** This conservation measure would
35 be implemented in the BDCP long-term implementation period because the design
36 of restored freshwater intertidal marshes would be dependent on the design and
37 construction of a new around-Delta conveyance channel.

38

39 **Implementation Considerations:** Implementation considerations include:

- 40 ▪ the feasibility for subsidence reversal using tule plantings or other
- 41 technique to raise ground surface elevations before breaching levees;

- 1 ▪ ensuring compatibility with flood control functions of north Delta levees
2 and channels (e.g., McCormack-Williamson Tract);
- 3 ▪ restoration effects on upstream and downstream flood risk;
- 4 ▪ the need to incorporate design features and management strategies to
5 preclude or minimize the establishment of non-native submerged aquatic
6 vegetation and other undesirable non-native species;
- 7 ▪ locating and designing levee breaches to maximize the development of
8 intertidal marsh and minimize hydrodynamic conditions that favor non-
9 native predatory fish;
- 10 ▪ the need to incorporate design features that will promote the natural
11 establishment of marsh-associated covered plant species;
- 12 ▪ consideration for the effects of restoration-induced dampening of the tidal
13 range on subsequent marsh restoration designs;
- 14 ▪ potential for increasing mercury methylation and resuspension of
15 contaminants;
- 16 ▪ compatibility with the footprint and facilities associated with an around-
17 Delta aqueduct;
- 18 ▪ determining appropriate allowable land uses and management activities on
19 transitional grasslands conserved to accommodate future sea level rise; and
- 20 ▪ securing fee title or easements and the protection of privately own lands
21 within the ROA.

22
23 **Resiliency to future changes:** This conservation measure is expected to be fairly
24 resilient to future changes in hydrology and sea levels. Conserving higher
25 elevation transitional grassland habitats along the margins of restored marsh will
26 provide sufficient lands to accommodate the upslope establishment of intertidal
27 marsh as sea level rises. If the alignment of an around-Delta aqueduct is upslope
28 of restored habitats, however, the area available for accommodating sea level rise
29 may be constrained.

30
31 **Uncertainties/risks:** Restoration of subtidal aquatic habitats could result in
32 infestation of non-native submerged aquatic vegetation and increase the
33 abundance of non-native predators or vulnerability of covered fish species to
34 predation. Methylation of mercury may occur in intertidal zones, making
35 methylmercury bioavailable to plants, fish, and wildlife in and downstream of
36 restored marshes (Alpers et al. 2006). Additionally, there could be a short-term
37 risk associated with mobilizing pesticides, herbicides, and other contaminants into
38 the Delta following initial introduction of tidal flow onto agricultural lands.

39
40 **Monitoring and adaptive management considerations:** Opportunities for
41 adaptive management are related to assessing the effectiveness of restored
42 marshes to develop as functional covered species habitats and to produce food and

1 organic carbon in support of food web processes. Results of monitoring the
2 development of early marsh restorations would help inform improvements in the
3 design and management of subsequent marsh restorations. Results of monitoring
4 early restoration projects could also be used to develop cost effective management
5 techniques, if needed, to control the establishment of non-native species in
6 restored marshes. Some of the monitoring considerations include:

- 7 ▪ type and extent of use by covered fishes;
- 8 ▪ extent of phytoplankton, zooplankton, and macroinvertebrate production in
9 marsh channels;
- 10 ▪ load of organic carbon, phytoplankton, zooplankton, and macroinvertebrates
11 produced in marshes and subsequently exported into the Delta;
- 12 ▪ extent of native vegetation relative to non-native vegetation on the marsh plain;
- 13 ▪ extent of native aquatic vegetation relative to non-native aquatic vegetation;
- 14 ▪ growth and survival of rearing Sacramento splittail and Chinook salmon in
15 shallow subtidal aquatic habitats;
- 16 ▪ the establishment of habitat conditions suitable for the natural establishment of
17 marsh-associated covered plant species; and
- 18 ▪ levels of mercury methylation and biological uptake.

19
20 **Reversibility:** This conservation measure would be difficult to reverse because it
21 would require construction of new levees to re-isolate restored habitat areas from
22 tidal flow and pumping to remove water from reclaimed habitat areas.

23
24 **Conservation Measure FIMA1.3: Restore a mosaic of [redacted] to [redacted] acres of intertidal**
25 **marsh and shallow subtidal aquatic habitat within the West Delta Restoration**
26 **Opportunity Area.** Restored freshwater intertidal marsh and shallow subtidal aquatic
27 habitats would be designed to support the physical and biological attributes described in
28 Attachment A. The mosaic of habitats would include at least [redacted] acres of freshwater
29 intertidal marsh habitat. Areas suitable for restoration include Decker Island, portions of
30 Sherman Island, Jersey Island, Bradford Island, Twitchell Island, and Brannon Island,
31 and along portions of the north bank of the Sacramento River where elevations and
32 substrates are suitable. The purpose of restoring intertidal marsh in the west Delta is to
33 provide a continuous corridor of habitat and food productivity linking current and future
34 restored habitat in the Cache Slough Complex with habitat in Suisun Marsh and Bay and
35 to provide intertidal marsh habitat within the anticipated future eastward position of the
36 low salinity zone with sea level rise.

37
38 Design elements of this conservation measure are anticipated to include:
39

- 1 ▪ placing fill material on shallowly subsided restoration sites to raise land
 - 2 surfaces to elevations suitable for restoration of intertidal marsh³;
 - 3 ▪ planting tules, or other techniques, to raise ground surface elevations suitable
 - 4 for intertidal marsh restoration on shallowly subsided portions of islands and
 - 5 breaching levees when target elevations are achieved;
 - 6 ▪ breaching and setting back levees to provide for tidal exchange with restored
 - 7 habitats; and
 - 8 ▪ excavating channels to initiate development of dendritic channel networks
 - 9 within restored marshes.
- 10
- 11 **Rationale:** Restoring freshwater intertidal marsh and shallow subtidal aquatic
- 12 habitats is expected to reduce the adverse effects of stressors related to food and
- 13 habitat availability for the covered species by:
- 14 ▪ increasing rearing habitat area for Chinook salmon, Sacramento splittail, and
 - 15 possibly steelhead (Healey 2001, Brown 2003);
 - 16 ▪ providing future habitat areas for delta smelt and longfin smelt within the
 - 17 anticipated eastward movement of the low salinity zone with sea level rise;
 - 18 ▪ increasing the production of food for rearing salmonids, splittail, and other
 - 19 species (Kjelson et al. 1982; Siegel 2007);
 - 20 ▪ increasing the availability and production of food in the western Delta and
 - 21 Suisun Bay by exporting organic material via tidal flow from the marsh plain
 - 22 and organic carbon, phytoplankton, zooplankton, and other organisms produced
 - 23 in intertidal channels into the Delta (Siegel 2007);
 - 24 ▪ locally providing areas of cool water refugia for delta smelt (C. Enright pers.
 - 25 comm.);
 - 26 ▪ increasing the extent of habitat available for colonization by Mason’s lilaepsis;
 - 27 and
 - 28 ▪ increasing the extent of habitat for California black rail and tricolored blackbird.
- 29 Lands within the West Delta ROA represent the only location to implement
- 30 intertidal marsh restorations within the anticipated future location of the low
- 31 salinity zone with sea level rise. A substantial proportion of the suitable
- 32 restoration sites in this area are in public ownership.
- 33
- 34 **Recommended Implementation Timeframe:** This conservation measure could
- 35 be initiated in the BDCP near-term implementation period and continue to be
- 36 implemented over the term of the BDCP as restoration opportunities are identified.
- 37

³ Sources of fill material could include dredge material from ongoing dredging operations and dredge spoils and sand deposits on Decker Island, Brannon Island, and other nearby suitable sites.

1 **Implementation Considerations:** Implementation considerations include:

- 2 ▪ the availability of suitable fill material and feasibility for subsidence reversal;
- 3 ▪ consideration for the effects of restoration-induced dampening of the tidal
- 4 range on subsequent marsh restoration designs and local tidal
- 5 hydrodynamics;
- 6 ▪ the need to design levees and provide elevations suitable to accommodate
- 7 future sea level rise;
- 8 ▪ locating and designing levee breaches to maximize the development of
- 9 intertidal marsh and minimize hydrodynamic conditions that favor non-native
- 10 predatory fish;
- 11 ▪ coordination with Delta levee programs to ensure that restored habitats are
- 12 protected from adverse effects that could be associated with future levee
- 13 failures;
- 14 ▪ determining the appropriate allowable land uses and management activities
- 15 on transitional grasslands conserved to accommodate future sea level rise;
- 16 ▪ the need to incorporate design features and management strategies to
- 17 preclude or minimize the establishment and abundance of undesirable non-
- 18 native species;
- 19 ▪ potential for increasing mercury methylation and resuspension of
- 20 contaminants;
- 21 ▪ the need to incorporate design features that will promote the natural
- 22 establishment of marsh-associated covered plant species; and
- 23 ▪ the likelihood for removal of food produced from restored intertidal marshes
- 24 by non-native clams.

25
26 **Resiliency to future changes:** The resiliency of this conservation measure to
27 accommodate future sea level rise is limited because of the extent of subsidence
28 in the west Delta. It is expected, however, that restoration designs would
29 incorporate elements that would provide land surface elevations sufficient to
30 accommodate the upslope establishment of marsh over time as sea level rises.

31
32 **Uncertainties/risks:** Restoration of subtidal aquatic habitats could result in
33 establishment of *Egeria* and other non-native plants that reduce the ecological
34 benefits for restored subtidal aquatic habitats to covered species. The abundance
35 of non-native predators and competitor abundance could increase and the ability
36 to control these species is uncertain. Methylation of mercury may occur in
37 intertidal zones, making methylmercury bioavailable to plants, fish, and wildlife
38 in and downstream of restored marshes (Alpers et al. 2006). Large scale levee
39 failures, in the central Delta could reduce species and ecosystem benefits
40 associated with restored marshes in the west Delta depending on the effects of
41 changed hydrodynamic conditions on tidal range and salinity gradients in the west

1 Delta. There could be a short-term risk associated with mobilizing pesticides,
2 herbicides, and other contaminants into the Delta following initial introduction of
3 tidal flow onto agricultural lands.
4

5 **Monitoring and adaptive management considerations:** Opportunities for
6 adaptive management are related to assessing the effectiveness of restored
7 marshes to develop as functional covered species habitats and to produce food and
8 organic carbon in support of food web processes. Results of monitoring the
9 development of early marsh restoration projects would help inform improvements
10 in the design and management of subsequent marsh restorations. Results of
11 monitoring early restorations could also be used to develop cost effective
12 management techniques, if needed, to control the establishment of non-native
13 species in restored marshes. Some of the monitoring considerations include:

- 14 ▪ type and extent of use by covered fishes;
- 15 ▪ extent of phytoplankton, zooplankton, and macroinvertebrate production in
16 marsh channels;
- 17 ▪ load of phytoplankton, zooplankton, and macroinvertebrates exported into
18 the Delta and Suisun Bay;
- 19 ▪ extent of food produced from restored habitats that are consumed by non-
20 native clams;
- 21 ▪ extent of native vegetation relative to non-native vegetation in the restored
22 marsh;
- 23 ▪ extent of native relative to non-native submerged aquatic vegetation;
- 24 ▪ effects of habitat restoration on salinity gradients in the west Delta;
- 25 ▪ levels of mercury methylation and biological uptake;
- 26 ▪ organic carbon production in restored marshes and export to the Delta and
27 Suisun Bay; and
- 28 ▪ growth and survival of rearing Sacramento splittail and Chinook salmon in
29 shallow subtidal aquatic habitats.

30
31 **Reversibility:** This conservation measure would be difficult to reverse because
32 reversing the measure would require construction of new levees to re-isolate
33 restored habitat areas from tidal flow.
34

35 **Conservation Measure FIMA1.4: Restore a mosaic of [redacted] to [redacted] acres of intertidal**
36 **marsh, shallow subtidal aquatic, and transitional grassland habitat within the South**
37 **Delta Restoration Opportunity Area.** Restored freshwater intertidal marsh and shallow
38 subtidal aquatic habitats would be designed to support the physical and biological
39 attributes described in Attachment A. The mosaic of habitats would include at least [redacted]
40 acres of freshwater intertidal marsh habitat. Suitable sites for restoring freshwater
41 intertidal marsh include Fabian Tract, Union Island, Middle Roberts Island, and Lower

1 Roberts Island. Sites selected for restoration would be depend on the location and design
2 of the selected conveyance pathway and operations for the through-Delta component of
3 the dual conveyance facility. Selected sites would be those that would provide
4 substantial species and ecosystem benefits with the selected through-Delta conveyance
5 configuration.

6
7 Design elements of this conservation measure could include:

- 8 ▪ planting tules or other techniques to raise currently subsided ground surface
9 elevations suitable for intertidal marsh restoration on shallowly subsided portions
10 of islands and breaching levees when target elevations are achieved;
- 11 ▪ scalping higher elevation portions of islands to provide fill for placement on
12 subsidied portions of islands to raise surface elevations;
- 13 ▪ breaching and setting back levees to provide for tidal exchange with restored
14 habitats;
- 15 ▪ constructing cross levees where appropriate to protect property and preclude
16 inundation of deeply subsided portions of islands;
- 17 ▪ locating and designing levee breaches to maximize the development of intertidal
18 marsh and minimize hydrodynamic conditions that favor non-native predatory
19 fish; and
- 20 ▪ excavating channels to initiate development of dendritic channel networks within
21 restored marshes.

22
23 **Rationale:** Restoring freshwater intertidal marsh and shallow subtidal aquatic
24 habitats is expected to reduce the adverse effects of stressors related to food
25 availability and habitat availability for the covered species by:

- 26 ▪ increasing rearing habitat area for Sacramento splittail, Chinook salmon
27 produced in the San Joaquin River and other eastside tributaries, and possibly
28 steelhead (Healey 2001, Brown 2003);
- 29 ▪ providing future habitat areas for delta smelt and longfin smelt with the
30 anticipated eastward movement of the low salinity zone with sea level rise;
- 31 ▪ increasing the production of food for rearing salmonids, splittail, and other
32 species (Kjelson et al. 1982; Siegel 2007);
- 33 ▪ increasing the availability and production of food in the Delta and Suisun Bay
34 by export from the south Delta of organic material via tidal flow from the new
35 marsh plain and organic carbon, phytoplankton, zooplankton, and other
36 organisms produced in new intertidal channels (Siegel 2007);
- 37 ▪ locally providing areas of cool water refugia for delta smelt (C. Enright pers.
38 comm.);
- 39 ▪ increasing the extent of habitat available for colonization by Mason's lilaeopsis;
40 and

- 1 ▪ increasing the extent of habitat for California black rail and tricolored blackbird.
2
3 Additionally, in conjunction with dual conveyance operations, marsh restoration in
4 the south Delta could expand the current distribution of delta smelt into formerly
5 occupied habitat areas.

6 **Recommended Implementation Timeframe:** This conservation measure would
7 need to be implemented following completion of the around-Delta facilities to
8 minimize adverse affects of through-Delta operations on restoration benefits.
9 Restoration planning, however, could be initiated in the near-term implementation
10 period.

11
12 **Implementation Considerations:** Implementation considerations include:

- 13 ▪ selecting the location for habitat restoration (Fabian Tract, Union Island,
14 Middle Roberts Island, or Lower Roberts Island) is dependent on the
15 through-Delta conveyance corridor (i.e., Old River or Middle River)
16 selected for dual operations and therefore the relative influence of South
17 Delta SWP/CVP pumping operations on the restored habitat;
- 18 ▪ feasibility of raising land surface elevations using tule plantings or other
19 techniques to raise ground surface elevations before breaching levees;
- 20 ▪ consideration of the effects of restoration-induced dampening of the tidal
21 range on local tidal hydrodynamics and subsequent marsh restoration
22 designs;
- 23 ▪ coordination with Delta levee programs to ensure that restored habitats are
24 protected from adverse effects that could be associated with future levee
25 failures;
- 26 ▪ locating and designing levee breaches to maximize the development of
27 intertidal marsh and minimize hydrodynamic conditions that favor non-native
28 predatory fish;
- 29 ▪ ensuring that designs for restored intertidal marshes along the San Joaquin
30 River would be compatible and provide synergistic species and ecosystem
31 benefits with proposed restoration of adjoining floodplain habitat upstream of
32 French Camp Slough as described under Conservation Measure FLOO2.1;
- 33 ▪ net level of species and ecosystem benefits that can be achieved with dual
34 conveyance operations;
- 35 ▪ potential for increasing mercury methylation and resuspension of
36 contaminants;
- 37 ▪ determining the appropriate allowable land uses and management activities
38 on transitional grasslands conserved to accommodate future sea level rise;
- 39 ▪ the need to incorporate design features and management strategies to
40 preclude or minimize the establishment and abundance of undesirable non-
41 native species;

- 1 ▪ the need to incorporate design features that will promote the natural
- 2 establishment of marsh-associated covered plant species; and
- 3 ▪ securing fee title or easements and the protection of privately own lands
- 4 within the ROA.

5

6 **Resiliency to future changes:** This conservation measure is expected to be fairly

7 resilient to future changes in hydrology and sea level. Conserving higher

8 elevation transitional grassland habitats along the margins of restored marsh will

9 provide sufficient lands to accommodate the upslope establishment of intertidal

10 marsh as sea level rises.

11

12 **Uncertainties/risks:** Restoration of subtidal aquatic habitats could result in

13 establishment of *Egeria* and other non-native plants that reduce the ecological

14 benefits of restored marsh for covered species. The abundance of non-native

15 predator and competitor abundance could increase and the ability to control them

16 is uncertain. Methylation of mercury may occur in intertidal zones, making

17 methylmercury bioavailable to plants, fish, and wildlife in and downstream of

18 restored marshes (Alpers et al. 2006). Large scale levee failures in the central

19 Delta could reduce species and ecosystem benefits associated with restored

20 marshes in the south Delta depending on the effects of changed hydrodynamic

21 conditions on tidal range and salinity gradients.

22

23 **Monitoring and adaptive management considerations:** Opportunities for

24 adaptive management are related to assessing the effectiveness of restored

25 marshes to develop as functional covered species habitats and to produce food and

26 organic carbon in support of food web processes. Results of monitoring the

27 development of early marsh restorations would help inform improvements in the

28 design and management of subsequent marsh restorations. Results of monitoring

29 early restorations could also be used to develop cost effective management

30 techniques, if needed, to control the establishment of non-native species in

31 restored marshes. Some of the monitoring considerations include:

- 32 ▪ type and extent of use by covered fishes;
- 33 ▪ extent of phytoplankton, zooplankton, and macroinvertebrate production in
- 34 marsh channels;
- 35 ▪ load of phytoplankton, zooplankton, and macroinvertebrates exported into
- 36 the central and west Delta;
- 37 ▪ organic carbon production in restored marshes and exported to the central
- 38 and west Delta;
- 39 ▪ levels of mercury methylation and biological uptake;
- 40 ▪ extent of native vegetation relative to non-native vegetation at marsh
- 41 surface;

- 1 ▪ effects of through-Delta operations on the amount of organic carbon and
- 2 food produced from restored marshes that is successfully exported to the
- 3 central and west Delta;
- 4 ▪ extent of native relative to non-native aquatic vegetation; and
- 5 ▪ growth and survival of rearing Sacramento splittail, Chinook salmon, and
- 6 other covered fish species in shallow subtidal aquatic habitats.

7
8 **Reversibility:** This conservation measure would be difficult to reverse because
9 reversal would require construction of new levees to re-isolate restored habitat
10 areas from tidal flow.

11
12 **Conservation Measure FIMA1.5: Restore a mosaic of [redacted] to [redacted] acres of intertidal**
13 **marsh, shallow subtidal aquatic, and transitional grassland habitat within the East**
14 **Delta Restoration Opportunity Area.** Restored freshwater intertidal marsh and shallow
15 subtidal aquatic habitats would be designed to support the physical and biological
16 attributes described in Attachment A. The mosaic of habitats would include at least [redacted]
17 acres of freshwater intertidal marsh habitat. Areas suitable for restoration include
18 Terminus Tract south of State Highway 12, Shin Kee Tract, Rio Blanco Tract, and
19 Bishop Bract. Design elements of this conservation measure could include:

- 20 ▪ acquisition of lands in fee-title or through conservation easements suitable for
- 21 restoration of intertidal and subtidal habitats and for accommodating future sea
- 22 level rise;
- 23 ▪ constructing levees to isolate deeply subsided lands and protect property;
- 24 ▪ planting tules or placing fill material to raise elevations of shallowly subsided
- 25 lands;
- 26 ▪ creating channels to promote the development of dendritic tidal channels; and
- 27 ▪ breaching levees to reintroduce tidal exchange to leveed lands.

28 If the eastern alignment of an around-Delta aqueduct is constructed, the aqueduct levee
29 (on its east side) may be incorporated into the design of restored marshes.

30
31 **Rationale:** Restoring freshwater intertidal marsh and shallow subtidal aquatic
32 habitats within the East Delta ROA is expected to reduce the adverse effects of
33 stressors related to food and habitat availability for the covered fish species by:

- 34 ▪ increasing rearing habitat area for Sacramento splittail and San Joaquin Chinook
- 35 salmon and possibly steelhead (Healey 2001, Brown 2003);
- 36 ▪ increasing the production of food for rearing salmonids, splittail, and other
- 37 species (Kjelson et al. 1982, Siegel 2007);
- 38 ▪ increasing the availability and production of food in the east and central Delta
- 39 by exporting organic material from the marsh plain and phytoplankton,
- 40 zooplankton, and other organisms produced in intertidal channels into the Delta
- 41 (Siegel 2007);

- 1 ▪ locally providing areas of cool water refugia for delta smelt (C. Enright pers.
2 comm.);
- 3 ▪ increasing the extent of habitat available for colonization by Mason's lilaeopsis,
4 and
- 5 ▪ increasing the extent of habitat for giant garter snake, California black rail, and
6 tricolored blackbird.

7
8 **Recommended Implementation Timeframe:** It is anticipated that this
9 conservation measure would be implemented in the BDCP long-term
10 implementation period because the design of restored freshwater intertidal marshes
11 would be influenced by the construction of a new around-Delta conveyance
12 facilities.

13
14 **Implementation Considerations:** Implementation considerations include:

- 15 ▪ the feasibility for subsidence reversal using tule plantings or other
16 techniques to raise ground surface elevations before breaching levees;
- 17 ▪ the need to incorporate design features and management strategies to
18 preclude or minimize the establishment of *Egeria* and other undesirable
19 non-native species;
- 20 ▪ locating and designing levee breaches to maximize the development of
21 intertidal marsh and minimize hydrodynamic conditions that favor non-
22 native predatory fish;
- 23 ▪ the need to incorporate design features that will promote the natural
24 establishment of marsh-associated covered plant species;
- 25 ▪ consideration for the effects of restoration-induced dampening of the tidal
26 range and local tidal hydrodynamics on subsequent marsh restoration
27 designs;
- 28 ▪ the footprint and facilities associated with an around-Delta aqueduct;
- 29 ▪ potential for increasing mercury methylation and resuspension of
30 contaminants;
- 31 ▪ determining the appropriate allowable land uses and management activities
32 on transitional grasslands conserved to accommodate future sea level rise;
33 and
- 34 ▪ securing fee-title or easements and the protection of privately own lands
35 within the ROA.

36 **Resiliency to future changes:** This conservation measure is expected to be fairly
37 resilient to future changes in hydrology and sea level. Conserving higher
38 elevation transitional grassland habitats along the margins of restored marsh
39 would provide lands to accommodate the upslope establishment of intertidal
40 marsh as sea level rises. If the alignment of an around-Delta aqueduct is upslope
41 of restored habitats, however, the area available for accommodating sea level rise

1 may be constrained.
2

3 **Uncertainties/risks:** Restoration of subtidal aquatic habitats could result in
4 infestation of non-native submerged aquatic vegetation and increase the
5 abundance of non-native predators and vulnerability of covered fish species to
6 predation. Methylation of mercury may occur in intertidal zones, making
7 methylmercury bioavailable to plants, fish, and wildlife in and downstream of
8 restored marshes (Alpers et al. 2006). Additionally, there could be a short-term
9 risk associated with mobilizing pesticides, herbicides, and other contaminants into
10 the Delta following initial introduction of tidal flow onto agricultural lands.
11

12 **Monitoring and adaptive management considerations:** Opportunities for
13 adaptive management are related to assessing the effectiveness of restored
14 marshes to develop as functional covered species habitats and to produce food and
15 organic carbon in support of food web processes. Results of monitoring the
16 development of early marsh restoration projects would help inform improvements
17 in the design and management of subsequent marsh restorations. Results of
18 monitoring early restorations could also be used to develop cost effective
19 management techniques, if needed, to control the establishment of non-native
20 species in restored marshes. Some of the monitoring considerations include:

- 21 ▪ type and extent of use by covered fishes;
- 22 ▪ extent of organic carbon, phytoplankton, zooplankton, and macroinvertebrate
23 production in marsh channels;
- 24 ▪ load of organic carbon, phytoplankton, zooplankton, and macroinvertebrates
25 exported into the Delta;
- 26 ▪ extent of native vegetation relative to non-native vegetation on the marsh plain;
- 27 ▪ extent of native submerged aquatic plants relative to non-native submerged
28 aquatic vegetation;
- 29 ▪ growth and survival of rearing Sacramento splittail, Chinook salmon, and other
30 covered fish species in shallow subtidal aquatic habitats;
- 31 ▪ the establishment of habitat conditions suitable for the natural establishment of
32 marsh-associated covered plant species; and
- 33 ▪ levels of mercury methylation and biological uptake.

34
35 **Reversibility:** This conservation measure would be difficult to reverse because it
36 would require construction of new levees to re-isolate restored habitat areas from
37 tidal flow and pumping to remove water from reclaimed habitat areas.
38

39 40 **Brackish Intertidal Marsh Habitat Restoration** 41 **Conservation Measures** 42

1 **Conservation Measure BIMA1.1 Restore a mosaic of [redacted] to [redacted] acres of brackish**
2 **intertidal marsh, shallow subtidal aquatic, and transitional grassland habitat within**
3 **the Suisun Marsh Restoration Opportunity Area.** Restored brackish intertidal marsh
4 would be designed to support the physical and biological attributes described in
5 Attachment A. The Suisun Marsh Restoration Plan (in development) currently provides
6 for restoring 6,000-9,000 acres of brackish intertidal marsh (S. Chappell pers. comm.).
7 Under this conservation measure, additional brackish intertidal marsh would be restored
8 opportunistically over the term of the BDCP as lands become available for restoration
9 from willing participants. Habitat would be restored as a mosaic of brackish intertidal
10 marsh, shallow subtidal aquatic, and transitional grassland habitats of which at least [redacted]
11 acres would be brackish intertidal marsh. Anticipated actions to restore brackish
12 intertidal marsh habitat include:

- 13 ▪ acquisition of lands in fee-title or through conservation easements suitable for
14 restoration of intertidal and subtidal habitats and for accommodating future sea
15 level rise from willing landowners;
- 16 ▪ planting tules or other techniques to raise elevations of shallowly subsided
17 lands;
- 18 ▪ reconnecting disconnected remnant sloughs to Suisun Bay and removing
19 remnant slough dikes to reintroduce tidal connectivity to slough watersheds to
20 restore tidal marsh; and
- 21 ▪ breaching dikes to reintroduce tidal exchange to diked lands.

22 Hydrodynamic modeling conducted for the Suisun Marsh Restoration Plan (J. DeGeorge
23 pers. comm.) indicates that restoring marsh north of Montezuma Slough would shift the
24 low salinity zone westward and restoring marsh at sites adjacent to Suisun Bay would
25 shift the low salinity zone eastward, potentially adversely affecting delta smelt habitat
26 and water quality in the west Delta. Consequently, implementation of marsh restoration
27 projects in north and south Suisun Marsh would be sequenced such that these potential
28 effects would be minimized.

29
30 **Rationale:** Restoring brackish intertidal marsh within Suisun Marsh is expected
31 to reduce the adverse effects of stressors related to food and habitat availability for
32 the covered species by:

- 33 ▪ increasing rearing habitat area for Chinook salmon, Sacramento splittail, and
34 possibly steelhead (Healey 2001, Siegel 2007);
- 35 ▪ increasing the production of food for rearing salmonids, splittail, and other
36 species (Kjelson et al. 1982);
- 37 ▪ increasing the availability and production of food in Suisun Bay by exporting
38 organic material via tidal flow from the marsh plain and phytoplankton,
39 zooplankton, and other organisms produced in intertidal channels into the Bay;
- 40 ▪ locally providing areas of cool water refugia for delta smelt (C. Enright pers.
41 comm.);

- 1 ▪ reducing periodic low dissolved oxygen events associated with the discharge of
- 2 waters from lands managed as seasonal freshwater wetlands that would be
- 3 restored as brackish intertidal marsh (Siegel 2007, C. Enright pers. comm.); and
- 4 ▪ increasing the extent of habitat available for colonization by Suisun marsh aster
- 5 and soft-bird's beak.

6 Additionally, the Suisun Marsh ROA encompasses a substantial area of land with

7 elevations suitable for intertidal marsh restoration that would involve few impacts

8 on infrastructure or permanent crops relative to the availability of suitable lands

9 within the Delta.

10

11 **Recommended Implementation Timeframe:** This conservation measure could

12 be initiated in the BDCP near-term implementation period and be implemented

13 over the term of the BDCP as restoration opportunities are identified.

14

15 **Implementation Considerations:** Implementation considerations include:

- 16 ▪ coordination with the Solano Multi-Species Habitat Conservation Plan and
- 17 the Suisun Marsh Plan to ensure effective implementation of conservation
- 18 measures among the plans;
- 19 ▪ feasibility for subsidence reversal using tule plantings or other techniques to
- 20 raise ground surface elevations before breaching levees;
- 21 ▪ consideration for the effects of restoration-induced dampening of the tidal
- 22 range and local tidal dynamics on subsequent marsh restoration designs;
- 23 ▪ the need to incorporate design features and management strategies to
- 24 preclude or minimize the establishment and abundance of undesirable non-
- 25 native species;
- 26 ▪ the need to incorporate design features that will promote the natural
- 27 establishment of marsh-associated covered plant species;
- 28 ▪ locating and designing levee breaches to maximize the development of
- 29 intertidal marsh and minimize hydrodynamic conditions that favor non-native
- 30 predatory fish;
- 31 ▪ evaluating the impact of likely removal of food produced from restored
- 32 brackish intertidal marshes by clams;
- 33 ▪ effects of operation of the salinity control gates on species and ecosystem
- 34 benefits provided by restored marshes;
- 35 ▪ potential for increasing mercury methylation and resuspension of
- 36 contaminants;
- 37 ▪ determining the appropriate allowable land uses and management activities
- 38 on transitional grasslands or managed seasonal wetlands conserved to
- 39 accommodate future sea level rise;
- 40 ▪ selecting restoration lands and implementing restoration in a sequence that

1 minimizes adverse effects of breaching/removing dikes on position of the low
2 salinity zone; and

- 3 ▪ securing fee-title or easements from willing private landowners and the
4 protection of privately lands within the ROA.

5
6 **Resiliency to future changes:** This conservation measure is expected to be fairly
7 resilient to future changes in hydrology and sea level. The landward margins of
8 Suisun Marsh border higher elevation transitional grassland habitats that would
9 provide sufficient lands for the upslope re-establishment of brackish intertidal
10 marsh as sea level rises and inundates marshes restored in those locations.

11 Sediment modeling conducted for existing proposed restorations in Suisun Marsh
12 also indicate that sediment supplies entering the marsh from tributaries may be
13 sufficient to allow the marsh plain south of Montezuma Slough to accrete at rates
14 that would keep pace with sea level rise (C. Enright, pers. comm.).

15
16 **Uncertainties/risks:** Restoration of subtidal aquatic habitats could result in
17 establishment of non-native plants that reduce the ecological benefits of restored
18 marsh for covered species. Non-native predator and competitor abundance could
19 increase and the ability to control them is uncertain. Initial studies have indicated
20 that sediment supplies are likely sufficient to allow for subsided lands south of
21 Montezuma Slough to accrete to form marsh plain. If restored habitats are
22 designed around this assumption and sediment supplies are not sufficient, restored
23 habitats would not provide the desired covered species benefits and could increase
24 the abundance of predators and competitors, adversely affecting covered fish
25 species. Altering existing habitat conditions in this area could potentially
26 adversely affect delta smelt habitat if salinity gradients, turbidity, or temperature
27 conditions change significantly as a result of restoration actions.

28
29 **Monitoring and adaptive management considerations:** Opportunities for
30 adaptive management are related to assessing the effectiveness of restored
31 marshes to develop as functional covered species habitats and to produce food and
32 organic carbon in support of food web processes. Results of monitoring the
33 development of early marsh restoration projects would help inform improvements
34 in the design and management of subsequent marsh restorations project. Results
35 of monitoring early restorations could also be used to develop cost effective
36 management techniques, if needed, to control the establishment of non-native
37 species in restored marshes. Some of the monitoring considerations include:

- 38 ▪ type and extent of use by covered fishes;
- 39 ▪ extent of organic carbon, phytoplankton, zooplankton, and
40 macroinvertebrate production in marsh channels;
- 41 ▪ load of organic carbon, phytoplankton, zooplankton, and
42 macroinvertebrates exported into Suisun Bay;
- 43 ▪ extent of food produced from restored habitats that are consumed by clams;

- 1 ▪ extent of native marsh vegetation relative to non-native vegetation;
- 2 ▪ effects of habitat restoration on salinity gradients and local tidal
- 3 hydrodynamics in the western Delta;
- 4 ▪ growth and survival of rearing Sacramento splittail, Chinook salmon, and
- 5 other covered species in shallow subtidal aquatic habitats.

6
7 **Reversibility:** This conservation measure would be difficult to reverse because
8 reversal would require construction of new dikes to re-isolate restored habitat
9 areas from tidal flow.

10 11 12 **Channel Margin Habitat Restoration Conservation Measures**

13
14 **Conservation Measures CHMA1.1. Support development and implementation of**
15 **levee construction and maintenance designs that incorporate aquatic and riparian**
16 **habitat features.** The BDCP Implementing Entity would coordinate with DWR, State
17 Reclamation Board, and U.S. Army Corps of Engineers to track planned levee
18 construction and maintenance activities. The BDCP Implementing Entity would
19 participate in planning processes for the construction of new levees, or maintenance of
20 existing levees, located along important habitat areas for covered fish species (e.g., fish
21 migration corridors). These activities will help ensure that levee designs incorporate
22 features that would benefit covered fish species, minimize adverse effects of the actions
23 on covered fish species, and avoid potential adverse effects of proposed actions on the
24 ecological functions provided by existing and planned BDCP conserved habitats.

25
26 **Rationale:** Improperly designed levees could increase habitat for non-native
27 predators, attract covered fish species, and thus contribute to increased predation
28 losses of covered fish species. Properly designed levees can support habitat for
29 salmonids and splittail. Riparian vegetation provides cover and rearing habitat for
30 covered fish species and organic carbon inputs into adjacent channels (U.S. Fish
31 and Wildlife Service 2004).

32
33 **Implementation timeframe:** This measure could be implemented in the BDCP
34 near-term implementation period and for the duration of the BDCP.

35
36 **Implementation Considerations:** Implementation considerations include
37 establishing a process that effectively engages the Implementing Entity in DWR,
38 Central Valley Flood Protection Board, and U.S. Army Corps of Engineers levee-
39 related planning processes.

40
41 **Resiliency to future changes:** If levees are sufficiently high and properly
42 designed to support vegetation, then riparian vegetation could move up the levee
43 face with the anticipated rising sea level.
44

1 **Uncertainties/risks:** There are uncertainties related to designing levee habitat
2 features that would improve habitat conditions for covered fish species and
3 degrade habitat conditions for non-native predatory fish. Restoring aquatic levee
4 habitats potentially increase the predation risk for covered fish species.
5

6 **Monitoring and adaptive management considerations:** It is anticipated that
7 lead agencies would include provisions for adaptive management and monitoring
8 in their levee planning documents. Adaptive management opportunities could
9 include monitoring the effectiveness of various levee habitat design components
10 and, based on monitoring results, adjusting levee habitat designs to improve
11 benefits for covered species. Some of the monitoring considerations include:

- 12 ▪ monitoring the use of aquatic levee habitats by covered fish species and
13 non-native predatory fish;
- 14 ▪ the natural establishment and regeneration of riparian vegetation on levee
15 slopes; and
- 16 ▪ the extent of zooplankton and macroinvertebrate production along restored
17 channel margin habitats compared to unvegetated levees.
18

19 **Reversibility:** Riparian habitat components of this measure are moderately
20 reversible as riparian vegetation established on or adjacent to levees could be
21 removed if necessary for levee repair, maintenance, or other reasons. Reversing
22 structural habitat design features (e.g., submerged low rock benches), however,
23 would be difficult.
24

25 **Conservation Measures CHMA1.2. Design levees constructed under the BDCP to**
26 **incorporate design features that support and enhance covered species habitats.**
27 BDCP site-specific habitat restoration designs may require the construction of new levees
28 (e.g., setback levees to restore floodplain habitat area). The BDCP Implementing Entity
29 would design such levees to incorporate design features that would provide for the
30 establishment of riparian and tidal emergent vegetation along low elevation surfaces (e.g.,
31 levee benches).
32

33 **Rationale:** Improperly designed levees could increase habitat for non-native
34 predators, attract covered fish species, and contribute to increased predation losses
35 of covered fish species. Properly designed levees can support and enhance habitat
36 for salmonids, splittail, and other covered fish species. Riparian vegetation
37 provides cover for covered fish species, and provides organic carbon inputs into
38 adjacent channels (U.S. Fish and Wildlife Service 2004).
39

40 **Implementation timeframe:** This measure could be implemented in the BDCP
41 near-term implementation period and for the duration of the BDCP.
42

43 **Implementation Considerations:** Implementation considerations include
44 coordinating with the U.S. Army Corps of Engineers, DWR, and other flood
45 control agencies to ensure that BDCP levee designs, as applicable, comply with

1 levee flood control standards.

2
3 **Resiliency to future changes:** If levees are sufficiently high and properly
4 designed to support vegetation, then riparian vegetation could move up the levee
5 face with the anticipated rising sea level.

6
7 **Uncertainties/risks:** There are uncertainties related to designing levee habitat
8 features that would improve habitat conditions for covered fish species and
9 degrade habitat conditions for non-native predatory fish. Restoring aquatic levee
10 habitats potentially increase the predation risk for covered fish species.

11
12 **Monitoring and adaptive management considerations:** Adaptive management
13 opportunities could include monitoring the effectiveness of various levee habitat
14 design components and, based on monitoring results, adjusting levee habitat
15 designs to improve benefits for covered species. Some of the monitoring
16 considerations include:

- 17 ▪ monitoring the use of aquatic levee habitats by covered fish species and
18 non-native predatory fish;
- 19 ▪ the natural establishment and regeneration of riparian vegetation on levee
20 slopes; and
- 21 ▪ the extent of zooplankton and macroinvertebrate production along restored
22 channel margin habitats compared to unvegetated levees.

23
24 **Reversibility:** Riparian habitat components of this measure are moderately
25 reversible as riparian vegetation established on or adjacent to levees could be
26 removed if necessary for levee repair, maintenance, or other reasons. Reversing
27 structural habitat design features (e.g., submerged low rock benches), however,
28 would be difficult.

29
30 **Conservation Measure CHMA1.3: Enhance channel margin habitats along [redacted] to [redacted]**
31 **miles of Steamboat and Sutter Sloughs to improve habitat conditions for covered**
32 **fish species.** Within the Steamboat Slough ROA, Steamboat and Sutter Sloughs are
33 thought to serve as important rearing habitat and movement corridors for juvenile
34 salmonids outmigrating from the Sacramento River (J. Burau pers. comm.). Habitat
35 conditions for covered fish species would be enhanced along [redacted] to [redacted] miles of Steamboat
36 Slough and [redacted] to [redacted] miles of Sutter Slough. The purpose of this measure is to improve
37 the growth and survival of juvenile salmonids that use these habitat areas.

38
39 Design elements for this conservation measure could include:

- 40 ▪ modifying channel geometry to improve hydrodynamic and structural
41 complexity for native species and to create conditions that are less favorable to
42 non-native fish predators and competitors; and
- 43 ▪ reducing the abundance of non-native fish predators and competitors.

1 Following implementation of habitat enhancements, the BDCP Implementing Entity
2 would undertake actions to encourage the transport of juvenile salmonids into Sutter
3 Slough if monitoring results indicate that survival and growth of juvenile salmonids that
4 rear and pass through Sutter and Steamboat Sloughs is substantially higher than under
5 current conditions. Increasing the proportion of juvenile salmonids transported into the
6 sloughs could be accomplished either by reorienting the upstream mouth of Sutter Slough
7 to the Sacramento River or constructing structures in the Sacramento River channel near
8 the upstream mouths of the sloughs that would guide the movement of fish into
9 Steamboat and Sutter Sloughs. To undertake this action, the BDCP Implementing Entity
10 would need to coordinate with and receive approvals from the U.S. Army Corps of
11 Engineers to either modify the project levees or construct an in-channel structure.

12
13 **Rationale:** Enhancing Steamboat and Sutter Sloughs as fish migration corridors
14 within the Steamboat Slough ROA is expected to increase the survival and growth
15 of outmigrating Sacramento River salmonids by:

- 16 ■ increasing the quality of rearing habitat area for Sacramento River salmonids (J.
17 Bureau pers. comm., Siegel 2007);
- 18 ■ reducing the risk for predation on covered fish species by non-native fish
19 predators (J. Bureau pers. comm.); and
- 20 ■ reducing the risk for entrainment of juvenile salmonids by providing a migration
21 corridor that bypasses the intakes of a new north Delta diversion point, the Delta
22 Cross Channel, and Georgiana Slough.

23
24 **Recommended Implementation Timeframe:** It is anticipated that this
25 conservation measure would be implemented in the BDCP long-term
26 implementation period.

27
28 **Implementation Considerations:** Implementation considerations include:

- 29 ■ the relative efficacy of various predatory fish control methods;
- 30 ■ appropriate modifications to the channel geometries of Steamboat and
31 Sutter Sloughs that could effectively improve habitat conditions for
32 juvenile salmonids and other species and degrade habitat conditions for
33 non-native predatory fish; and
- 34 ■ coordination with the U.S. Army Corps of Engineers and other flood
35 control agencies to allow for: 1) modifications to project levees or
36 placement of in-channel structures to improve transport of juvenile
37 salmonids into Steamboat and Sutter Sloughs and 2) modifications to the
38 channel geometry of the sloughs.

39
40 **Resiliency to future changes:** This conservation measure is expected to be fairly
41 resilient to future changes in hydrology and sea levels because the types of habitat
42 improvements are such that they would be expected to continue to provide greater
43 benefits for juvenile salmonids than under future conditions without the

1 improvements.

2
3 **Uncertainties/risks:** The efficacy of the proposed restoration actions for
4 increasing survival and growth of juvenile salmonids by reducing predation risk is
5 uncertain, particularly if flow velocities are substantially reduced as a result of
6 increasing flows into the Yolo Bypass and operating a new Delta diversion.

7
8 **Monitoring and adaptive management considerations:** Opportunities for
9 adaptive management are related to assessing the effectiveness of restoration
10 actions in improving the survival and growth of juvenile salmonids passing
11 through the sloughs by improving habitat conditions and reducing predation and
12 entrainment risk. Results of monitoring could help inform the BDCP
13 Implementing Entity of subsequent opportunities to improve these sloughs as
14 salmonid rearing habitats and migration corridors. Implementation of this
15 conservation measure would also afford the opportunity to test fish predator
16 control techniques to identify the most efficacious methods for controlling
17 predator populations. Some of the monitoring considerations include assessing
18 the:

- 19 ▪ change in survival and growth of juvenile salmonids using the sloughs relative
20 to current conditions;
- 21 ▪ effectiveness of channel geometry designs for improving salmonid rearing
22 habitat and degrading non-native predatory fish habitat;
- 23 ▪ effectiveness of predatory fish control methods; and
- 24 ▪ effectiveness of channel modifications for increasing the transport of juvenile
25 salmonids into the sloughs.

26 **Reversibility:** This conservation measure could be difficult to reverse depending
27 on the magnitude and nature of channel modifications.

30 **Riparian Habitat Restoration Conservation Measures**

31
32 **Conservation Measure RIPA1.1. Restore between [redacted] and [redacted] acres of riparian forest**
33 **and scrub communities as a component of restored floodplain, freshwater intertidal**
34 **marsh, and channel margin habitats.** As described in Attachment A, the design of
35 restored floodplain, freshwater intertidal marsh, and channel margin habitats [see
36 Conservation Measures FLOO 1.1, FLOO2.1, FLOO 2.2, FIMA1.1-1.5, BIMA1.1,
37 CHMA1.1 and 1.2] will incorporate restoration of riparian habitats as described below.

38
39 **Floodplain Habitat Restoration.** To the extent consistent with flood control
40 requirements, restored floodplain habitat areas will allow for the natural establishment
41 and growth of woody riparian vegetation on portions of restored floodplains that support
42 appropriate soils and hydrology. At floodplain restoration sites that function
43 hydrologically as flood bypasses (e.g., the Yolo Bypass), riparian vegetation is expected

1 to establish along margins of existing and created drains and channels and other locations
2 with suitable hydrology. In bypasses co-managed for habitat and flood control benefits,
3 locations where riparian vegetation is allowed to establish would be limited to areas
4 where the presence of riparian vegetation would not compromise flood control standards
5 or hydraulic capacity of the flood control bypass.

6
7 Riparian habitat would be allowed to naturally establish in floodplain habitat areas that
8 are restored by setting back levees to expand the extent of the floodplain subject to
9 overbank flow.

10
11 **Freshwater Intertidal Marsh Restoration.** Woody riparian vegetation will be allowed
12 to naturally reestablish along the upper elevation margins of restored intertidal marsh
13 habitats where soils and hydrology are suitable, including segments of stream channels
14 that drain into restored marshes.

15
16 **Channel Margin Habitat Restoration.** As described under Conservation Measure
17 CHMA1.2, BDCP levees will be designed to provide for the establishment and growth of
18 riparian vegetation along levees. Levees constructed and maintained by other entities
19 that incorporate “green” levee components would also increase the extent of riparian
20 habitat within the Planning Area by allowing for the establishment and growth of riparian
21 vegetation on levee surfaces.

22
23 **Rationale:** Restoring riparian forest and riparian scrub habitats is expected to
24 provide the following ecosystem and covered species benefits:

- 25
- 26 ■ increasing the extent of valley elderberry longhorn beetle habitat and nesting
27 habitat for Swainson’s hawk and yellow breasted chat;
 - 28 ■ increasing the extent of shaded riverine aquatic cover and increasing instream
29 cover by through contributions of instream woody material (U.S. Fish and
30 Wildlife Service 2004);
 - 31 ■ providing inputs of organic material (e.g., leave and twig drop) in support of
32 aquatic foodweb processes; and
 - 33 ■ increasing cover for rearing juvenile salmonids and Sacramento splittail.

34
35 **Recommended Implementation Timeframe:** It is anticipated that elements of
36 this conservation measure would be implemented in both near-term and long-term
37 BDCP implementation period.

38
39 **Implementation Considerations:** Implementation considerations include
40 ensuring that designs for the floodplain, intertidal marsh, and channel margin
41 habitat restorations described under Conservation Measures FLOO 1.1, FLOO2.1,
42 FLOO 2.2, FIMA1.1-1.6, BIMA1.1, CHMA1.1 and 1.2 provide for the restoration
43 of at least [redacted] acres of riparian forest and scrub habitat and the potential need for
44 periodic control of non-native invasive plant species. Other implementation

1 considerations for this conservation measure are included under implementation
2 considerations for Conservation Measures FLOO 1.1, FLOO2.1, FLOO 2.2,
3 FIMA1.1-1.5, BIMA1.1, CHMA1.1 and 1.2.
4

5 **Resiliency to future changes:** Restored riparian habitats are expected to be
6 fairly resilient to future changes in hydrology and sea level rise because habitats
7 will be restored within large sites that would be expected to provide a sufficient
8 range of site characteristics (e.g., elevation and soil gradients) to allow for the
9 ongoing reestablishment of riparian vegetation in response to changes in
10 hydrologic and sea level conditions over time.
11

12 **Uncertainties/risks:** Allowing for the natural establishment of native riparian
13 vegetation could result in the establishment of riparian habitats dominated by non-
14 native invasive species.
15

16 **Monitoring and adaptive management considerations:** Opportunities for
17 adaptive management include improving the design and management of restored
18 floodplain, channel margin, and freshwater intertidal marsh to provide for the
19 successfully establishment, growth, and benefits of restored riparian habitats
20 based on monitoring of the development of previously restored riparian habitats.
21 For example, if the natural establishment and growth of native riparian vegetation
22 is substantially impaired by competition with non-native plants, restoration
23 projects may need to provide for the control of non-native plants or require that
24 riparian plantings be installed to improve restoration success. Some of the
25 monitoring considerations include assessing the:

- 26 ■ use of restored riparian habitats by valley elderberry longhorn beetle,
27 Swainson's hawk, yellow-breasted chat, and riparian brush rabbit;
- 28 ■ factors governing the natural establishment and growth of native riparian
29 vegetation over a range of site conditions associated with restored floodplain,
30 channel margin, and intertidal marsh habitat areas;
- 31 ■ the need to control non-native plants to provide for the natural establishment of
32 native riparian vegetation; and
- 33 ■ ability for native riparian vegetation to reestablish in patterns that provide
34 desired ecosystem and covered species benefits.

35 **Reversibility:** The reversibility of riparian habitat restorations are the same as
36 described for each of the ROA restoration actions described under Conservation
37 Measures FLOO 1.1, FLOO2.1, FLOO 2.2, FIMA1.1-1.5, BIMA1.1, CHMA1.1
38 and 1.2.
39

40 References

41 Printed References

42
43

- 1 Alpers CN, C Eagles-Smith, C Foe, S Klasing, MC Marvin-DiPasquale, DG Slotton, L
2 Windham-Myers. 2008. Mercury conceptual model. Sacramento(CA): Delta
3 Regional Ecosystem Restoration Implementation Plan.
4
- 5 Brown L.R. 2003. Will Tidal Wetland Restoration Enhance Populations of Native
6 Fishes? In: Larry R. Brown, editor. Issues in San Francisco Estuary Tidal
7 Wetlands Restoration. San Francisco Estuary and Watershed Science. Vol. 1,
8 Issue 1 (October 2003), Article 2.
9 <http://repositories.cdlib.org/jmie/sfews/vol1/iss1/art2>
10
- 11 Feyrer, F., T. Sommer, and W. Harrell. 2006. Managing floodplain inundation for native
12 fish: production dynamics of age-0 splittail (*Pogonichthys macrolepidotus*) in
13 California's Yolo Bypass. *Hydrobiologia* 573:213-226.
14
- 15 Healey M.C. 1991. Life history of Chinook salmon (*Oncorhynchus tshawytscha*). In:
16 Groot C, Margolis L, editors. Pacific salmon life histories. Vancouver (BC):
17 University of British Columbia Press. p 311-394.
18
- 19 Jassby A. 2008. Phytoplankton in the upper San Francisco Estuary: recent biomass
20 trends, their causes and their trophic significance. San Francisco Estuary and
21 Watershed Science. Vol. 6, Issue 1 (February), Article 2.
22
- 23 Kjelson MA, Raquel PF, Fisher FW. 1982. Life history of fall-run juvenile chinook
24 salmon, *Oncorhynchus tshawytscha*, in the Sacramento-San Joaquin Estuary,
25 California. In: Kennedy VS, editor. Estuarine comparisons. New York (NY):
26 Academic Press. p 393-411.
27
- 28 Mitsch, W. J., J.G. Gosselink. 2000. Wetlands 3rd ed. Wiley & Sons Inc., New York.
29
- 30 Moss, B. 2007. Ecology of Fresh Waters 3rd ed. Blackwell Publishing, Oxford.
31
- 32 Moyle, P. B. 2002. Inland Fishes of California, 2nd edition. University of California Press,
33 Berkeley, California.
34
- 35 Moyle, P. B., R. D. Baxter, T. Sommer, T. C. Foin, and S. A. Matern. 2004. Biology and
36 population dynamics of the Sacramento splittail (*Pogonichthys macrolepidotus*) in
37 the San Francisco Estuary: a review. *San Francisco Estuary and Watershed*
38 *Science* 2(2):1-47.
39
- 40 Siegel S. W. 2007. Starting the Discussion: Foundation concepts and some initial
41 activities to restore ecosystem functions to the California Delta.
42
- 43 Sommer T, Harrell B, Nobriga M, Brown R, Moyle P, Kimmerer W, Schemel L. 2001a.
44 California's Yolo Bypass: evidence that flood control can be compatible with
45 fisheries, wetlands, wildlife, and agriculture. *Fisheries* 26(8):6-16.
46

1 Sommer, T. R., M. L. Nobriga, W. C. Harrell, W. Batham, and W. J. Kimmerer. 2001b.
2 Floodplain rearing of juvenile Chinook salmon: Evidence of enhanced growth and
3 survival. *Canadian Journal of Fisheries and Aquatic Sciences* 58(2):325-333.

4
5 Sommer, T. R., L. Conrad, G. O'Leary, F. Feyrer, and W. C. Harrell. 2002. Spawning and
6 rearing of splittail in a model floodplain wetland. *Transactions of the American*
7 *Fisheries Society* 131(5):966-974.

8
9 Sommer, T. R., R. D. Baxter, and F. Feyrer. 2007. Splittail “delisting”: A review of
10 recent population trends and restoration activities. *American Fisheries Society*
11 *Symposium* 53:25-38.

12
13 Sommer T.R., W.C. Harrell, M. Zoltan, F. Feyrer. 2008. Habitat associations and
14 behavior of adult and juvenile splittail (*Cyprinidae: Pogonichthys*
15 *macrolepidotus*) in a managed seasonal floodplain wetland. *San Francisco Estuary*
16 *and Watershed Science* 6(2):1-16.

17
18 USFWS. 2004. Impacts of riprapping to aquatic organisms and river functioning, Lower
19 Sacramento River, California. 2nd Edition. Available at:
20 www.fws.gov/sacramento/es/animal_spp_account/acctfish.htm.

21 22 23 **Personal Communications**

24
25 J. Burau, Engineer, US Geological Survey, Conversation with Hood Bypass Subgroup,
26 August 28, 2008.

27
28 S. Chappell, Executive Director, Suisun Resource Conservation District, personal
29 meeting, April 25, 2008.

30
31 J. DeGeorge, Vice President, RMA, Presentation to the Habitat and Operations Technical
32 Team, May 25, 2008.

33
34 C. Enright, Senior Water Resources Engineer, DWR, Presentation to the Habitat and
35 Operations Technical Team on May 28, 2008 and in DRERIP evaluation session
36 conducted on July 28, 2008.

Attachment A. Restoration Concepts for Habitats

Definitions of Restoration Concepts

This attachment describes the floodplain, intertidal marsh, and channel margin restoration concepts developed by the Habitat Restoration Program Technical Team (HRPTT).

These descriptions are intended to provide guidance to the BDCP Implementing Entity for planning habitat restoration actions and to initially define the range of physical and biological conditions that must be present in restored habitat areas in order for the restoration to be considered successful. The draft information presented in the restoration concept descriptions will be developed further and incorporated into the BDCP Conservation Strategy chapter. Each description includes the following information:

Restoration Variables: Brief descriptions of the key physical parameters that can be manipulated through restoration design and operations to restore habitat under the concept.

Design Targets: Narrative description of the desired physical and biological conditions that are expected to develop in restored habitat areas as a result of manipulating restoration variables.

Desired Ecological Benefits: Brief descriptions of covered fish species stressor effects expected to be reduced with implementation of the restoration concept.

Potential Performance Criteria (monitoring needs and adaptive management triggers): Physical and biological parameters that can be measured and that are indicators of the extent of desired ecological functions to be provided by habitats restored under the concept. The performance criteria represent the range of indicators that may be appropriate to monitor to assess the effectiveness of restored habitats in achieving desired covered species and ecosystem benefits. Results of monitoring may be used to trigger adaptive management responses through the BDCP adaptive management process to improve the effectiveness of restored habitats to provide desired benefits.

Key Uncertainties: Brief descriptions of major unknowns with respect to designing habitat restorations and benefits that are expected to be afforded by restoration habitats.

Potential Ecological Risks: Brief descriptions of potential unintended adverse physical and biological impacts that could be associated with implementing the restoration concept.

Floodplain Restoration Concept

Restoration Variables

- Seasonal timing of inundation
- Interannual frequency of inundation
- Spatial extent of inundation
- Depth of inundation
- Water velocity
- Connectivity with intertidal marsh and open water habitats
- Accessibility to migrating fish
- Design related to stranding risk and fish passage
- Vegetation type and cover
- Dry season land use (compatible farming practices)
- Grading/slope

Design Targets

Inundated Floodplain

- Shallow with highly variable depth (2 feet deep on average)
- Adequate hydraulic residence time to promote primary and secondary food production and export and turbidity export (number of days to produce desired food resources)
- Average velocities of about 1.5 foot/sec, but highly variable spatially and temporally
- Duration of inundation about 30-45 days
- Relatively large area (>1,000 acres) to accrue substantive benefit to fish populations
- Stranding avoided through good drainage
- Provides for passage around weirs or other inflow control structures
- Minimized risk for problem levels of methyl mercury and other contaminants
- Inundated during periods that favor native fish and disfavor non-native fish predators – generally late winter to early-mid spring
- Hydrodynamic variability through floodplain cross-section via heterogeneous topography

- 1 ▪ Flows exit floodplain via a channel system that, where possible, flows through
- 2 intertidal marsh towards open water
- 3 ▪ Natural connectivity to adjacent uplands to provide transitional habitats and
- 4 accommodate species movement

5

6 **Dry Floodplain**

7

- 8 ▪ Minimized use of persistent pesticides that are toxic to aquatic organisms
- 9 ▪ Cover and type of residual standing crop biomass (for floodplains with flood
- 10 protection function) or riparian and perennial vegetation (for floodplains without
- 11 flood protection function)
- 12 ▪ Allow for the natural establishment of woody riparian vegetation to the extent
- 13 consistent with desired land uses and flood control requirements

14

15 **Desired Ecological Benefits**

16

- 17 ▪ Primary and secondary production
- 18 ▪ Primary and secondary production export to Delta
- 19 ▪ Export of allochthonous material to Delta
- 20 ▪ Substantial increase in high quality splittail spawning and rearing habitat and
- 21 Chinook salmon (all runs) and steelhead rearing habitat relative to existing in-
- 22 Delta habitat conditions
- 23 ▪ Reduction in stranding/poaching losses of adult sturgeon and salmonids below
- 24 Fremont Weir
- 25 ▪ Improved habitat connectivity between upstream and downstream habitats
- 26 ▪ Improved survival/escapement of juvenile salmonids
- 27 ▪ Improved turbidity conditions (?)

28

29 **Potential Performance Criteria (possible monitoring needs and adaptive**

30 **management triggers)**

31

- 32 ▪ Extent of phytoplankton/zooplankton/macroinvertebrate production on
- 33 floodplain
- 34 ▪ Extent of phytoplankton/zooplankton/macroinvertebrate exported to the Delta
- 35 ▪ Growth rate of juvenile salmonids on floodplains
- 36 ▪ Proportion of outmigrating juvenile salmonids accessing floodplain habitats (by
- 37 run)
- 38 ▪ Extent of splittail spawning

- 1 ▪ Extent of native fish stranding
- 2 ▪ Extent of successful upstream passage of adult salmonids and sturgeon
- 3 ▪ Extent of mercury methylation
- 4 ▪ Contaminant load exported to Delta
- 5 ▪ Extent of habitat connectivity along migratory routes for anadromous fishes

6 7 **Key Uncertainties** 8

- 9 ▪ Proper depth for optimizing fish habitat conditions and food production
- 10 ▪ Proper inundation duration/residence time for optimizing fish growth and
11 survival and food production
- 12 ▪ Conditions necessary for the natural establishment of channel-associated
13 covered plant species in floodplains restored by setting back levees
- 14 ▪ Benefits of floodplain inundation to sturgeon, particularly juveniles, are
15 undocumented

16 17 **Potential Ecological Risks**

- 18 ▪ Mercury methylation
- 19 ▪ Establishment of non-native invasive species into created habitat

20 21 22 **Freshwater Intertidal Marsh Restoration Concept** 23

24 **Restoration Variables** 25

- 26 ▪ Spatial distribution of restored habitats within the Delta
- 27 ▪ Extent, location, and configuration of restored habitat
- 28 ▪ Amplitude of tidal exchange
- 29 ▪ Size and location of levee breaches
- 30 ▪ Channel cross sectional profile (elevation of marsh plain, topographic diversity,
31 depth, and slope)
- 32 ▪ Intertidal marsh channel density

33 34 **Design Targets** 35

- 36 ▪ Dominated by native freshwater emergent vegetation (predominantly tules,)

- 1 ▪ Presence of sinuous, dendritic channel networks of high density
- 2 ▪ Sufficient tidal exchange to promote primary and secondary production and its
- 3 export into the aquatic food web
- 4 ▪ Located throughout the Delta for optimal use by and benefit to covered species
- 5 ▪ Located where it can filter non-point source pollution from surface or subsurface
- 6 infiltration
- 7 ▪ High velocity, shallow channels to potentially prevent establishment of non-
- 8 native submerged aquatic vegetation that supports non-native predator habitat
- 9 ▪ Large tidal connectivity to open water areas to minimize steep flow velocity
- 10 gradients that promote establishment of non-native submerged aquatic
- 11 vegetation and provide predatory fish habitats
- 12 ▪ Natural connectivity to adjacent uplands to provide transitional habitats and
- 13 accommodate species movement
- 14 ▪ Accessible to fish, but does not trap fish
- 15 ▪ Connectivity with other intertidal marshes and with floodplain, open water,
- 16 channel margin, and low gradient upland habitats
- 17 ▪ Located such that other stressors (e.g., diversions) do not substantially reduce
- 18 functions beneficial to covered species
- 19 ▪ Designed to allow localized reductions in water temperature though nocturnal
- 20 thermal reduction

21 22 **Desired Ecological Benefits**

- 23
- 24 ▪ Primary and secondary production
- 25 ▪ Primary and secondary production export to Delta channels
- 26 ▪ Reduced summer/fall water temperature through nocturnal thermal exchange
- 27 and reintroduction of cooled water to Delta waterways
- 28 ▪ Filter for contaminants or site for transformation of contaminants
- 29 ▪ Splittail and salmonid rearing habitat
- 30 ▪ Potential delta smelt, longfin smelt, and splittail spawning habitat

31 32 **Potential Performance Criteria (possible monitoring needs and adaptive**

33 **management triggers)**

- 34
- 35 ▪ Type and extent of use by covered fishes
- 36 ▪ Extent of in-marsh phytoplankton/zooplankton/macroinvertebrate production
- 37 ▪ Extent of phytoplankton/zooplankton/macroinvertebrate exported into the Delta

- 1 ▪ Extent of native vegetation relative to non-native vegetation at marsh surface
- 2 ▪ Extent of native relative to non-native submerged aquatic vegetation
- 3 ▪ Extent of organic carbon production and export to Delta channels

4 5 **Key Uncertainties** 6

- 7 ▪ Ability to control non-native submerged aquatic vegetation and fish
- 8 ▪ Ability to restore native plant species (e.g., Delta tule pea)
- 9 ▪ Availability of adequate sediment supply and rate of tule growth for marsh
10 accretion
- 11 ▪ Extent and effectiveness for providing aquatic covered species and ecosystem
12 benefits
- 13 ▪ Effects of increased dampening of the tidal range as marsh restorations are
14 implemented on the ability to implement subsequent restorations
- 15 ▪ Effect of freshwater tidal marsh restoration on water quality and hydrodynamics
16 upstream and downstream

17 18 **Potential Ecological Risks** 19

- 20 ▪ Possibility of establishment of non-native invasive species into restored habitats
- 21 ▪ Depending on location, benefits may be reduced by diversions (project and non-
22 project)

23 24 25 **Brackish Intertidal Marsh Restoration** 26

27 **Restoration Variables** 28

- 29 ▪ Extent, location, and configuration of restored habitat
- 30 ▪ Distribution along salinity gradient
- 31 ▪ Amplitude of tidal exchange
- 32 ▪ Delta freshwater outflow
- 33 ▪ Size and location of dike breaches
- 34 ▪ Channel cross sectional profile (elevation of marsh plain, topographic diversity,
35 depth, and slope)
- 36 ▪ Intertidal marsh channel density

37

1 Design Targets

2

- 3 ▪ Dominated by native brackish marsh vegetation (e.g., pickleweed, saltgrass)
- 4 ▪ Presence of sinuous, dendritic channel networks of high density
- 5 ▪ Adjacent to higher elevation uplands to accommodate future with sea level rise
- 6 ▪ Primarily low marsh
- 7 ▪ Sufficient tidal exchange to promote primary and secondary production and its
- 8 export into the estuarine food web
- 9 ▪ Natural connectivity to adjacent uplands to provide transitional habitats and
- 10 accommodate species movement
- 11 ▪ Restore habitats that provide a range of salinity gradients
- 12 ▪ Accessible to fish, but does not trap fish
- 13 ▪ Connectivity with other intertidal marshes and with floodplain, open water,
- 14 channel margin, and upland habitats
- 15 ▪ Located such that other stressors (e.g., diversions) do not substantially reduce
- 16 functions beneficial to covered species
- 17 ▪ Designed to allow localized reductions in water temperature though nocturnal
- 18 thermal reduction

19

20 Desired Ecological Benefits

21

- 22 ▪ Primary and secondary production
- 23 ▪ Primary and secondary production export to Suisun Bay
- 24 ▪ Reduced summer/fall water temperature through nocturnal thermal exchange
- 25 and reintroduction of cooled water to Delta waterways
- 26 ▪ Filter for contaminants or site for transformation of contaminants
- 27 ▪ Splittail, salmonid, and sturgeon rearing habitat

28

29 Potential Performance Criteria (possible monitoring needs and adaptive 30 management triggers)

31

- 32 ▪ Type and extent of use by covered fishes
- 33 ▪ Extent of in-marsh phytoplankton/zooplankton/macroinvertebrate production
- 34 ▪ Extent of phytoplankton/zooplankton/macroinvertebrate exported into Suisun
- 35 Bay
- 36 ▪ Extent of native vegetation relative to non-native vegetation at marsh surface

- 1 ▪ Extent of organic carbon production and export into Suisun Bay

2

3 **Key Uncertainties**

4

- 5 ▪ Ability to control non-native fish (e.g., inland silversides)
- 6 ▪ Ability to restore native plant species (Suisun Marsh aster and soft bird's-beak)
- 7 ▪ Availability of adequate sediment supply for marsh accretion
- 8 ▪ Extent and effectiveness for providing aquatic covered species and ecosystem
- 9 benefits
- 10 ▪ Effects of increased dampening of the tidal range as marsh restorations are
- 11 implemented on the ability to implement subsequent restorations
- 12 ▪ Effect of brackish tidal marsh restoration on the position of the low salinity zone

13

14 **Potential Ecological Risks**

15

- 16 ▪ Possibility of establishment of non-native invasive species into restored habitat

17

18

19 **Channel Margin Habitat Restoration Concept**

20

21 **Restoration Variables**

22

- 23 ▪ Spatial distribution, extent, and location within the Delta
- 24 ▪ Length of restored habitat along channel margins
- 25 ▪ Cross sectional profile (elevation of habitat, topographic diversity, width,
- 26 variability in edge and bench surfaces, depth, and slope)
- 27 ▪ Amount and distribution of installed large woody debris
- 28 ▪ Extent of shaded riverine aquatic cover and vegetation needed to provide future
- 29 inputs of large woody debris

30

31 **Design Targets**

32

- 33 ▪ Incorporate large woody debris in banks (i.e., complex structure refugia)
- 34 ▪ Provide range of hydrodynamic conditions to benefit natives and minimize the
- 35 colonization of non-native submerged aquatic vegetation and predators
- 36 ▪ Provide woody riparian vegetation to create overhead cover and refuge from
- 37 predators in roots
- 38 ▪ Located and configured to connect to existing patches of habitat

- 1 ▪ Minimize use by predatory fish
- 2 ▪ Minimize occurrence of non-native submerged aquatic vegetation
- 3 ▪ Located along fish movement corridors and rearing habitats

4 5 **Desired Ecological Benefits**

- 6
- 7 ▪ Improved local and diurnal water temperatures at a local scale
- 8 ▪ Splittail spawning habitat
- 9 ▪ Splittail and salmonid rearing habitat
- 10 ▪ Source of allochthonous material
- 11 ▪ Phytoplankton/zooplankton/macrobenthos production
- 12 ▪ Increased hydrodynamic complexity in channels

13 14 **Potential Performance Criteria (possible monitoring needs and adaptive**

15 **management triggers)**

- 16
- 17 ▪ Type and extent of use by covered fishes
- 18 ▪ Type and extent of use by non-native predatory fish
- 19 ▪ Extent of overhead cover and woody riparian vegetation
- 20 ▪ Extent of native vegetation relative to non-native vegetation
- 21 ▪ Extent of phytoplankton/zooplankton/macrobenthos production

22 23 **Key Uncertainties**

- 24
- 25 ▪ Cost:benefit ratio associated with improving channel margin habitats along
- 26 levees

27 28 **Potential Ecological Risks**

- 29
- 30 ▪ Possibility of establishment of non-native invasive species into created habitat